arXiv:math/0501398v1 [math.AG] 23 Jan 2005

Opers.

by A. Beilinson and V. Drinfeld

In this article, which we wrote for ourselves in 1993, there are no new results, we only rewrote some definitions from [DS1] and [DS2] in a coordinate-free manner. We did this exercise because of the applications of opers to the geometric Langalnds program.

We decided to publish this text as an e-print because it can be used by students.

©§1. *G*-opers and $\mathfrak{g}$-opers.

©.1.1 Let G be a connected reductive group over $\mathbb{C}$ with a fixed Borel subgroup $B \subset G$ and a fixed Cartan subgroup $H \subset B$. Let $\mathfrak{g} \supset \mathfrak{b} \supset \mathfrak{h}$ be the corresponding Lie algebras, $\Gamma \subset \mathfrak{h}^*$ the set of simple roots with respect to $\mathfrak{b}$. If $\alpha \in \mathfrak{h}^*$ we set $\mathfrak{g}^\alpha = \{x \in \mathfrak{g} | [a,x] = \alpha(a)x$ for all $a \in \mathfrak{h}$'s. There is a unique Lie algebra grading $\mathfrak{g} = \underset{k}{\oplus}\mathfrak{g}_k$ such that $\mathfrak{g}_0 = \mathfrak{h}$, $\mathfrak{g}_1 = \underset{\alpha\in\Gamma}{\oplus}\mathfrak{g}^\alpha$ and $\mathfrak{g}_{-1} = \underset{\alpha\in\Gamma}{\oplus}\mathfrak{g}^{-\alpha}$. The corresponding filtration $\mathfrak{g}^k = \underset{r\geq k}{\oplus}\mathfrak{g}_r$ is B-invariant.

Fix a smooth algebraic curve Y over $\mathbb{C}$. Given a B-bundle $\mathcal{F}$ on Y we denote by $\mathcal{E}_\mathcal{F}$ the algebroid of infinitesimal symmetries of $\mathcal{F}$ and by $\mathcal{E}^\mathfrak{g}_\mathcal{F}$ the algebroid of infinitesimal symmetries of the G-bundle corresponding to $\mathcal{F}$. We have a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathfrak{b}_\mathcal{F} & \longrightarrow & \mathcal{E}_\mathcal{F} & \longrightarrow & \Theta & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \| & & \\ 0 & \longrightarrow & \mathfrak{g}_\mathcal{F} & \longrightarrow & \mathcal{E}^\mathfrak{g}_\mathcal{F} & \longrightarrow & \Theta & \longrightarrow & 0 \end{array} \qquad (©.1)$$

with exact rows where $\mathfrak{b}_\mathcal{F}$ and $\mathfrak{g}_\mathcal{F}$ denote the $\mathcal{F}$-twists of $\mathfrak{b}$ and $\mathfrak{g}$ while Θ denotes the tangent bundle of Y. We have the filtration $\mathfrak{g}^k_\mathcal{F} \subset \mathfrak{g}_\mathcal{F}$ and also the filtration $\mathcal{E}^k_\mathcal{F} \subset \mathcal{E}^\mathfrak{g}_\mathcal{F}, k \leq 0$, where $\mathcal{E}^k_\mathcal{F}$ denotes the preimage of $\mathfrak{g}^k_\mathcal{F}/\mathfrak{b}_\mathcal{F} \subset \mathfrak{g}_\mathcal{F}/\mathfrak{b}_\mathcal{F} = \mathcal{E}^\mathfrak{g}_\mathcal{F}/\mathcal{E}_\mathcal{F}$. Notice that $\mathcal{E}^{-1}_\mathcal{F}/\mathcal{E}_\mathcal{F} = \mathfrak{g}^{-1}_\mathcal{F}/\mathfrak{g}_\mathcal{F} = \underset{\alpha\in\Gamma}{\oplus}\mathfrak{g}^{-\alpha}_\mathcal{F}$. Here $\mathfrak{g}^{-\alpha}_\mathcal{F}$ denotes the $\mathcal{F}$-twist of the B-module $\mathfrak{g}^{-\alpha}$ (the action of B on $\mathfrak{g}^{-\alpha}$ is defined to be the composition $B \to H \to \operatorname{Aut}\mathfrak{g}^{-\alpha}$).

©.1.2 Definition. *A G-oper on Y is a B-bundle $\mathcal{F}$ on Y with a connection $\nabla : \Theta \to \mathcal{E}^\mathfrak{g}_\mathcal{F}$ such that $\nabla(\Theta) \subset \mathcal{E}^{-1}_\mathcal{F}$ and for every $\alpha \in \Gamma$ the composition*

$$\Theta \xrightarrow{\nabla} \mathcal{E}^{-1}_\mathcal{F} \longrightarrow \mathcal{E}^{-1}_\mathcal{F}/\mathcal{E}_\mathcal{F} \longrightarrow \mathfrak{g}^{-\alpha}_\mathcal{F}$$

is an isomorphism.

There is a natural notion of isomorphism of G-opers. So G-opers on Y form a groupoid.

Remark. The term "oper" is motivated by the fact that for most of the classical G one can interpret G-opers as differential operators between certain line bundles (see ©§2).

©.1.3 Let $\mathfrak{g}$ be a semisimple Lie algebra over $\mathbb{C}$.

Definition. *A $\mathfrak{g}$-oper is a G-oper where G is the group of inner automorphisms of $\mathfrak{g}$.*

Remark. The notion of $\mathfrak{g}$-oper (but not the word "oper") was introduced in [DS1] and [DS2] under the assumption that Θ is trivial and a trivialization of Θ is fixed. The notion of $sl(n)$-oper for an arbitrary Y was introduced in [Te1]. The notion of $sl(2)$-oper is equivalent to that of Sturm-Liouville operator or projective connection (see ©.2.6 and ©.2.7). So $sl(2)$-opers have a long history (see the survey [Tyu]).

Proposition. *A $\mathfrak{g}$-oper does not have nontrivial automorphisms.*

Proof. The assertion is local, so we may assume that the B-bundle $\mathcal{F}$ corresponding to our $\mathfrak{g}$-oper is trivial and that there is a regular function z on Y such that dz has no zeros. The connection ∇ on $\mathcal{F}$ is defined by an operator of the form $\partial_z + q$ where q is a regular function on Y with values in $\mathfrak{g}^{-1}$ such that for every $\alpha \in \Gamma$ the component $q_{-\alpha}$ of q has no zeros. An automorphism of $(\mathcal{F}, \nabla)$ is a B-valued regular function b on Y such that

$$b^{-1}\,(\partial_z + q)\,b = \partial_z + q \qquad (©.2)$$

Let q_k be the components of q with respect to the grading $\mathfrak{g} = \underset{k}{\oplus}\mathfrak{g}_k$. Then (©.2) implies that $b_0^{-1} q_{-1} b_0 = q_{-1}$ where b_0 is the image of b in H. Since the center of G is trivial $b_0 = 1$. So we may consider the logarithm $\log b$. Let u_k be the components of $\log b$ with respect to the grading $\mathfrak{g} = \underset{k}{\oplus}\mathfrak{g}_k$. Obviously $u_0 = 0$. Assume that $r > 0$ and $u_k = 0$ for $k < r$. Then it follows from (©.2) that $[q_{-1}, u_r] = 0$. Notice that q_{-1} is a member of an $sl(2)$-triple $\{h, x, y\}$ with $y = q_{-1}$ and $h \in \mathfrak{h}$ being the sum of fundamental coweights. Since $[h, u_r] = r u_r$ and $r > 0$ the equality $[y, u_r] = 0$ implies that $u_r = 0$. So $u_k = 0$ for all k and therefore $b = 1$. ■

Corollary. *If G is an arbitrary connected reductive group and Y is connected then the group of automorphisms of any G-oper on Y is equal to the center of G.* ■

©.1.4 Let G and $\mathfrak{g}$ denote the same objects as in ©.1.1. Denote by G_{ss} the commutant of G (i.e., the semisimple part of G) and by $\mathfrak{g}_{ss}$ the Lie algebra of G_{ss}. Let us discuss the relation between G-opers and $\mathfrak{g}_{ss}$-opers. Denote by Z the center of G and by Ω the canonical bundle of X. Of course to each G-oper there corresponds a (G/Z)-oper, i.e., a $\mathfrak{g}_{ss}$-oper. On the other hand if one fixes a square root of Ω (i.e., a line bundle $\mathcal{M}$ on X with an isomorphism $\mathcal{M}^{\otimes 2}\xrightarrow{\sim}\Omega$) then there is a natural way to lift a $\mathfrak{g}_{ss}$-oper to a G_{ss}-oper and therefore to a G-oper. Indeed, set $B_{\mathrm{ad}} = B/Z$, $H_{\mathrm{ad}} = H/Z$, $B_{ss} = B\cap G_{ss}$, $H_{ss} = H\cap G_{ss}$. Let $(\mathcal{F}_{\mathrm{ad}}, \nabla)$ be a $\mathfrak{g}_{ss}$-oper. $\mathcal{F}_{\mathrm{ad}}$ is a B_{ad}-bundle and lifting $(\mathcal{F}_{\mathrm{ad}}, \nabla)$ to a G_{ss}-oper means lifting $\mathcal{F}_{\mathrm{ad}}$ to a B_{ss}-bundle. Denote by $\mathcal{F}^H_{\mathrm{ad}}$ the H_{ad}-bundle corresponding to $\mathcal{F}_{\mathrm{ad}}$. Lifting $\mathcal{F}_{\mathrm{ad}}$ to a B_{ss}-bundle means lifting $\mathcal{F}^H_{\mathrm{ad}}$ to a H_{ss}-bundle. But for every $\alpha\in\Gamma$ we have a canonical isomorphism $\mathfrak{g}^{-\alpha}_{\mathcal{F}_{\mathrm{ad}}}\xrightarrow{\sim}\Theta$ (cf. ©.1.1) and therefore $\mathfrak{g}^{\alpha}_{\mathcal{F}_{\mathrm{ad}}}\xrightarrow{\sim}\Omega$. Fix non-zero elements $y_\alpha\in\mathfrak{g}^{-\alpha}$. Then the isomorphisms $\mathfrak{g}^{-\alpha}_{\mathcal{F}_{\mathrm{ad}}}\xrightarrow{\sim}\Theta$ induce an isomorphism between $\mathcal{F}^H_{\mathrm{ad}}$ and the direct image of the $\mathbb{G}_m$-torsor on Y corresponding to Ω with respect to the homomorphism $\lambda:\mathbb{G}_m\to H_{\mathrm{ad}}$ such that $\lambda(t)$ acts on each $\mathfrak{g}^\alpha, \alpha\in\Gamma$, as multiplication by t. Notice that there is a unique homomorphism $\mu:\mathbb{G}_m\to H_{ss}$ such that the diagram

$$\begin{array}{ccc}\mathbb{G}_m & \xrightarrow{t\mapsto t^2} & \mathbb{G}_m\\ {\scriptstyle\mu}\downarrow & & \downarrow{\scriptstyle\lambda}\\ H_{ss} & \longrightarrow & H_{\mathrm{ad}}\end{array} \qquad (©.3)$$

is commutative (indeed, λ corresponds to $\check\rho$, the sum of the fundamental coweights of $\mathfrak{g}_{ss}$, and since $2\check\rho$ belongs to the coroot lattice $2\check\rho$ corresponds to a homomorphism $\mathbb{G}_m\to H_{ss}$). So once $\Omega^{\otimes(1/2)}$ and y_α are fixed there is a natural way of lifting $\mathcal{F}^H_{\mathrm{ad}}$ to a H_{ss}-bundle: just take the direct image with respect to μ of the $\mathbb{G}_m$-torsor corresponding to $\Omega^{\otimes(1/2)}$. The dependence of the lifting on the choice of y_α is not very serious: the H_{ss}-bundles corresponding to $\{y_\alpha\}$ and $\{\widetilde{y}_\alpha\}$ $\big($and to the same choice of $\Omega^{\otimes(1/2)}\big)$ are isomorphic but the isomorphism depends on the choice of $h\in H_{ss}$ such that $hy_\alpha h^{-1}=\widetilde{y}_\alpha$ for all α.

So for a fixed $\Omega^{\otimes(1/2)}$ we have an equivalence between the category of G-opers and the category of pairs $(\mathcal{E}_1,\mathcal{E}_2)$ where $\mathcal{E}_1$ is a $\mathfrak{g}_{ss}$-oper and $\mathcal{E}_2$ is a Z-bundle with a connection: given $\mathcal{E}_1$ one constructs a G_{ss}-oper as explained above and then one twists it by means of $\mathcal{E}_2$. This equivalence depends on the choice of $\Omega^{\otimes(1/2)}$ in the following way. Let $\mathcal{M}$ and

$\widetilde{\mathcal{M}}$ be the square roots of Ω. If the pair $(\mathcal{E}_1, \mathcal{E}_2)$ corresponds to $\mathcal{M}$ and a certain G-oper and $(\widetilde{\mathcal{E}}_1, \widetilde{\mathcal{E}}_2)$ corresponds to $\widetilde{\mathcal{M}}$ and the same G-oper then

$$\widetilde{\mathcal{E}}_1 = \mathcal{E}_1,\ \widetilde{\mathcal{E}}_2 = \mathcal{E}_2 \otimes \nu_*\big(\widetilde{\mathcal{M}} \otimes \mathcal{M}^{\otimes(-1)}\big) \qquad (©.4)$$

Here $\widetilde{\mathcal{M}} \otimes \mathcal{M}^{\otimes(-1)}$ is considered as a torsor over the group $\{1, -1\}$ and $\nu_*\big(\widetilde{\mathcal{M}} \otimes \mathcal{M}^{\otimes(-1)}\big)$ is its image with respect to the homomorphism $\nu : \{1, -1\} \to Z$ such that $\nu(-1) = \mu(-1)$ $\big($the commutativity of (©.3) implies that $\mu(-1) \in \mathrm{Ker}(H_{ss} \to H_{ad}) = H_{ss} \cap Z\big)$.

©§2. G-opers for classical G.

We are going to show that for most of the classical groups G a G-oper can be considered as a differential operator between certain line bundles.

©.2.1 Let us start with the case $G = GL(n)$. In this case a G-oper can be considered as a locally free $\mathcal{O}_Y$-module sheaf of rank n with a filtration $\mathcal{E} = \mathcal{E}_n \supset \mathcal{E}_{n-1} \supset \ldots \supset \mathcal{E}_1 \supset \mathcal{E}_0 = 0$ and a connection $\nabla : \mathcal{E} \to \mathcal{E} \otimes \Omega$ such that

(i) the sheaves $\mathcal{E}_i/\mathcal{E}_{i-1}, 1 \leq i \leq n$, are invertible,

(ii) for $i = 1, 2, \ldots, n-1$ $\nabla(\mathcal{E}_i) \subset \mathcal{E}_{i+1} \otimes \Omega$ and ∇ induces an isomorphism $\mathcal{E}_i/\mathcal{E}_{i-1} \xrightarrow{\sim} (\mathcal{E}_{i+1}/\mathcal{E}_i) \otimes \Omega$.

Denote by $\mathcal{D}$ the sheaf of differential operators on Y and write $\mathcal{O}$ instead of $\mathcal{O}_Y$. Let $\mathcal{D}_i \subset \mathcal{D}$ be the sheaf of differential operators of order $\leq i$. The connection ∇ defines a $\mathcal{D}$-module structure on $\mathcal{E}$. So we have a natural morphism $\mathcal{D} \otimes_{\mathcal{O}} \mathcal{E}_1 \to \mathcal{E}$. The condition (ii) implies that the morphism $\mathcal{D}_{n-1} \otimes_{\mathcal{O}} \mathcal{E}_1 \to \mathcal{E}$ is an isomorphism. So considering the diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{D}_{n-1} \otimes_{\mathcal{O}} \mathcal{E}_1 & \longrightarrow & \mathcal{D}_n \otimes_{\mathcal{O}} \mathcal{E}_1 & \longrightarrow & \Omega^{\otimes(-n)} \otimes_{\mathcal{O}} \mathcal{E}_1 & \longrightarrow & 0 \\ & & & \searrow^{\sim} & \downarrow & & & & \\ & & & & \mathcal{E} & & & & \end{array}$$

we obtain a splitting $\Omega^{\otimes(-n)} \otimes_{\mathcal{O}} \mathcal{E}_1 \to \mathcal{D}_n \otimes_{\mathcal{O}} \mathcal{E}_1$ and therefore a section $L \in H^0\big(Y, \mathcal{B} \otimes_{\mathcal{O}} \mathcal{D} \otimes_{\mathcal{O}} \mathcal{A}^{\otimes(-1)}\big)$ where $\mathcal{A} = \mathcal{E}_1^{\otimes(-1)}$, $\mathcal{B} = \Omega^{\otimes n} \otimes \mathcal{E}_1^{\otimes(-1)}$. Consider L as a differential operator $\mathcal{A} \to \mathcal{B}$. Its order equals to n and its principal symbol $\sigma \in H^0\big(Y, \mathcal{A}^{\otimes(-1)} \otimes \mathcal{B} \otimes \Omega^{\otimes(-n)}\big)$ equals to 1.

Conversely, suppose we are given invertible sheaves $\mathcal{A}, \mathcal{B}$ and a differential operator $L : \mathcal{A} \to \mathcal{B}$ of order n whose principal symbol $\sigma \in H^0\big(Y, \mathcal{A}^{\otimes(-1)} \otimes \mathcal{B} \otimes \Omega^{\otimes(-n)}\big)$ has no zeros (in this situation σ defines an isomorphism $\mathcal{B} \xrightarrow{\sim} \mathcal{A} \otimes \Omega^{\otimes n}$ and after identifying $\mathcal{B}$ with $\mathcal{A} \otimes \Omega^{\otimes n}$ we may say that $\sigma = 1$). Then we construct a $GL(n)$-oper as follows. We have a $\mathcal{D}$-module morphism $r_L : \mathcal{D} \otimes_{\mathcal{O}} \mathcal{B}^{\otimes(-1)} \to \mathcal{D} \otimes_{\mathcal{O}} \mathcal{A}^{\otimes(-1)}$ defined by $r_L(u) = uL$ where L is considered as a section of $\mathcal{B} \otimes_{\mathcal{O}} \mathcal{D} \otimes_{\mathcal{O}} \mathcal{A}^{\otimes(-1)}$. Set $\mathcal{E} = \operatorname{Coker} r_L$, $\mathcal{E}_i = \operatorname{Im}\big(\mathcal{D}_i \otimes_{\mathcal{O}} \mathcal{A}^{\otimes(-1)} \to \mathcal{E}\big)$ and consider the $\mathcal{D}$-module $\mathcal{E}$ as an $\mathcal{O}$-module with a connection ∇. Then $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla)$ is an $GL(n)$-oper.

So we have constructed an *equivalence between the category of $GL(n)$-opers and the category of triples $(\mathcal{A}, \mathcal{B}, L)$ where $\mathcal{A}$ and $\mathcal{B}$ are invertible sheaves on Y and $L : \mathcal{A} \to \mathcal{B}$ is a differential operator of order n with non-vanishing principle symbol.* Let us explain that by a morphism from $(\mathcal{A}_1, \mathcal{B}_1, L_1)$ to $(\mathcal{A}_2, \mathcal{B}_2, L_2)$ we mean a pair of isomorphisms $\mathcal{A} \xrightarrow{\sim} \mathcal{A}_1$, $\mathcal{B} \xrightarrow{\sim} \mathcal{B}_1$ such that the diagram

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{L} & \mathcal{B} \\ \downarrow\wr & & \downarrow\wr \\ \mathcal{A}_1 & \xrightarrow{L_1} & \mathcal{B}_1 \end{array}$$

is commutative.

We can make a slightly more precise statement. Denote by $\operatorname{Diff}_n(\mathcal{A}, \mathcal{B})$ the set of differential operators $L : \mathcal{A} \to \mathcal{B}$ of order n with non-vanishing principal symbol. If $L \in \operatorname{Diff}_n(\mathcal{A}, \mathcal{B})$ and $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla)$ is the corresponding $GL(n)$-oper then we have canonical isomorphisms

$$\mathcal{E}_i/\mathcal{E}_{i-1} = \mathcal{E}_i \otimes \Omega^{\otimes(1-i)} = \mathcal{A}^{\otimes(-1)} \otimes \Omega^{\otimes(1-i)} = \mathcal{B}^{\otimes(-1)} \otimes \Omega^{\otimes(n+1-i)} \qquad (\text{©}.5)$$

In particular $\mathcal{E}_1 = \mathcal{A}^{\otimes(-1)}$, $\mathcal{E}/\mathcal{E}_{n-1} = \Omega \otimes \mathcal{B}^{\otimes(-1)}$. Denote by $\operatorname{Op}_n(\mathcal{A}, \mathcal{B})$ the set of isomorphism classes of $GL(n)$-opers $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla)$ with $\mathcal{E}_1 = \mathcal{A}^{\otimes(-1)}$ and $\mathcal{E}/\mathcal{E}_{n-1} = \Omega \otimes \mathcal{B}^{\otimes(-1)}$ (i.e., isomorphisms $\mathcal{E}_1 \xrightarrow{\sim} \mathcal{A}^{\otimes(-1)}$ and $\mathcal{E}/\mathcal{E}_{n-1} = \Omega \otimes \mathcal{B}^{\otimes(-1)}$ are fixed). We have constructed a canonical bijection

$$\operatorname{Op}_n(\mathcal{A}, \mathcal{B}) \xrightarrow{\sim} \operatorname{Diff}_n(\mathcal{A}, \mathcal{B}) \qquad (\text{©}.6)$$

©.2.2 Recall that the transpose L^t of a differential operator $L = \sum_i f_i \partial_z^i$ is defined by $L^t = \sum_i (-\partial_z)^i f_i$. In the global situation if L is a differential operator $\mathcal{A} \to \mathcal{B}$ where $\mathcal{A}$

and $\mathcal{B}$ are locally free $\mathcal{O}$-modules then L^t is a differential operator $\Omega\otimes\mathcal{B}^*\to\Omega\otimes\mathcal{A}^*$. In particular transposition maps $\mathcal{D}$ to $\Omega\otimes_{\mathcal{O}}\mathcal{D}\otimes_{\mathcal{O}}\Omega^{\otimes(-1)}$, i.e., the sheaf of differential operators $\Omega\to\Omega$. The induced morphism $\mathcal{A}\otimes_{\mathcal{O}}\mathcal{D}\otimes_{\mathcal{O}}\mathcal{B}\to\mathcal{B}\otimes\Omega\otimes_{\mathcal{O}}\mathcal{D}\otimes_{\mathcal{O}}\Omega^{\otimes(-1)}\otimes\mathcal{A}$ will also be called transposition and denoted by $L\mapsto L^t$.

If $\mathcal{A}$ and $\mathcal{B}$ are invertible sheaves on Y and $L\in\mathrm{Diff}_n(\mathcal{A},\mathcal{B})$ then $L^t\in\mathrm{Diff}_n\bigl(\Omega\otimes\mathcal{B}^{\otimes(-1)},\Omega\otimes\mathcal{A}^{\otimes(-1)}\bigr)$. On the other hand if $(\mathcal{E},\{\mathcal{E}_i\},\nabla)$ is a $GL(n)$-oper with $\mathcal{E}_1=\mathcal{A}^{\otimes(-1)}$ and $\mathcal{E}/\mathcal{E}_{n-1}=\Omega\otimes\mathcal{B}^{\otimes(-1)}$ then we can consider the dual $GL(n)$-oper $(\mathcal{E}^*,\{(\mathcal{E}^*)_i\},\nabla)$ where $(\mathcal{E}^*,\nabla)$ is the dual of $(\mathcal{E},\nabla)$ and $(\mathcal{E}^*)_i=\mathcal{E}_{n-i}^{\perp}\subset\mathcal{E}^*$. Since $(\mathcal{E}^*)_1=(\mathcal{E}/\mathcal{E}_{n-1})^*=\Omega^{\otimes(-1)}\otimes\mathcal{B}$ and $\mathcal{E}^*/(\mathcal{E}^*)_{n-1}=(\mathcal{E}_1)^*=\mathcal{A}$ dualization defines a mapping $\mathrm{Op}_n(\mathcal{A},\mathcal{B})\to\mathrm{Op}_n\bigl(\Omega\otimes\mathcal{B}^{\otimes(-1)},\Omega\otimes\mathcal{A}^{\otimes(-1)}\bigr)$.

Proposition. *The diagram*

$$\begin{array}{ccc}\mathrm{Diff}_n(\mathcal{A},\mathcal{B}) & \xrightarrow{L\mapsto -L^t} & \mathrm{Diff}_n\bigl(\Omega\otimes\mathcal{B}^{\otimes(-1)},\Omega\otimes\mathcal{A}^{\otimes(-1)}\bigr)\\ \downarrow\wr & & \downarrow\wr\\ \mathrm{Op}_n(\mathcal{A},\mathcal{B}) & \longrightarrow & \mathrm{Op}_n\bigl(\Omega\otimes\mathcal{B}^{\otimes(-1)},\Omega\otimes\mathcal{A}^{\otimes(-1)}\bigr)\end{array}$$

is commutative.

Proof. Let $(\mathcal{E},\{\mathcal{E}_i\},\nabla)$ and $(\widetilde{\mathcal{E}},\{\widetilde{\mathcal{E}}_i\},\widetilde{\nabla})$ be the $GL(n)$-opers corresponding respectively to L and $-L^t$. We must construct a horizontal non-degenerate pairing $\mathcal{E}\otimes\widetilde{\mathcal{E}}\to\mathcal{O}$ such that $\widetilde{\mathcal{E}}_i=\mathcal{E}_{n-i}^{\perp}$ and the isomorphisms $\mathcal{E}/\mathcal{E}_{n-1}\xrightarrow{\sim}\widetilde{\mathcal{E}}_1^*$ and $\widetilde{\mathcal{E}}/\widetilde{\mathcal{E}}_{n-1}\xrightarrow{\sim}\mathcal{E}_1^*$ induced by the pairing are equal to the compositions $\mathcal{E}/\mathcal{E}_{n-1}\xrightarrow{\sim}\Omega\otimes\mathcal{B}^{\otimes(-1)}\xrightarrow{\sim}\widetilde{\mathcal{E}}_1^*$ and $\widetilde{\mathcal{E}}/\widetilde{\mathcal{E}}_{n-1}\xrightarrow{\sim}\mathcal{A}\xrightarrow{\sim}\mathcal{E}_1^*$. More precisely the latter condition means the commutativity of the diagrams

$$\begin{array}{ccc}\mathcal{E}/\mathcal{E}_{n-1}=\mathcal{E}_1\otimes\Omega^{\otimes(1-n)} & = & \mathcal{A}^{\otimes(-1)}\otimes\Omega^{\otimes(1-n)}\\ \searrow^{\sim} & & \downarrow\varphi\\ \widetilde{\mathcal{E}}_1^* & = & \mathcal{B}^{\otimes(-1)}\otimes\Omega\end{array}\qquad(\text{©}.7\,\text{a})$$

$$\begin{array}{ccc}\widetilde{\mathcal{E}}/\widetilde{\mathcal{E}}_{n-1}=\widetilde{\mathcal{E}}_1\otimes\Omega^{\otimes(1-n)} & = & \mathcal{B}\otimes\Omega^{\otimes(-n)}\\ \searrow^{\sim} & & \downarrow(-1)^{n-1}\psi\\ \mathcal{E}_1^* & = & \mathcal{A}\end{array}\qquad(\text{©}.7\,\text{b})$$

where the equalities denote the canonical isomorphisms (©.4) while φ and ψ denote division by $\sigma\in H^0\bigl(Y,\mathcal{A}^{\otimes(-1)}\otimes\mathcal{B}\otimes\Omega^{\otimes(-n)}\bigr)$, the principle symbol of L. Let us explain that $(-1)^{n-1}$ appears in (©.7 b) because the principal symbol of $-L^t$ equals $(-1)^{n-1}\sigma$.

To construct the pairing we use the sheaf $\underline{\mathcal{D}}$ of pseudodifferential symbols $\big($local sections of $\underline{\mathcal{D}}$ can be written as formal expressions $\sum\limits_{i=-\infty}^{k} f_i\partial_z^i$ or as $\sum\limits_{i=-\infty}^{k} \partial_z^i g_i$ where f_i and g_i are regular functions and z is a local parameter on $Y\big)$. We also use the canonical morphism $\mathrm{res} : \underline{\mathcal{D}} \otimes_{\mathcal{O}} \Omega^{\otimes(-1)} \to \mathcal{O}$ defined by $\big\{\sum\limits_i f_i\partial_z^i\big\}(dz)^{-1} \mapsto f_{-1}$.

Here is the construction. We have $\mathcal{E} = \mathrm{Coker}\, r_L$ where $r_L : \mathcal{D} \otimes_{\mathcal{O}} \mathcal{B}^{\otimes(-1)} \to \mathcal{D} \otimes_{\mathcal{O}} \mathcal{A}^{\otimes(-1)}$ is defined by $r_L(u) = uL$, $L \in H^0\big(Y, \mathcal{B} \otimes_{\mathcal{O}} \mathcal{D} \otimes_{\mathcal{O}} \mathcal{A}^{\otimes(-1)}\big)$. We also have $\widetilde{\mathcal{E}} = \mathrm{Coker}\, r_{L^t}$, $r_{L^t} : \mathcal{D} \otimes_{\mathcal{O}} \mathcal{A} \otimes \Omega^{\otimes(-1)} \to \mathcal{D} \otimes_{\mathcal{O}} \mathcal{B} \otimes \Omega^{\otimes(-1)}$, $L^t \in H^0\big(Y, \Omega \otimes \mathcal{A}^{\otimes(-1)} \otimes_{\mathcal{O}} \mathcal{D} \otimes_{\mathcal{O}} \mathcal{B} \otimes \Omega^{\otimes(-1)}\big)$. Define a pairing between $\mathcal{D} \otimes_{\mathcal{O}} \mathcal{A}^{\otimes(-1)}$ and $\mathcal{D} \otimes_{\mathcal{O}} \mathcal{B} \otimes \Omega^{\otimes(-1)}$ by

$$(u, v) = \mathrm{res}\big(uL^{-1}v^t\big) \qquad (\text{©}.8)$$

Here $u \in \mathcal{D} \otimes_{\mathcal{O}} \mathcal{A}^{\otimes(-1)}$ $\big($i.e., u is a local section of $\mathcal{D} \otimes_{\mathcal{O}} \mathcal{A}^{\otimes(-1)}\big)$, $v \in \mathcal{D} \otimes_{\mathcal{O}} \mathcal{B} \otimes \Omega^{\otimes(-1)}$, $v^t \in \mathcal{B} \otimes_{\mathcal{O}} \mathcal{D} \otimes_{\mathcal{O}} \Omega^{\otimes(-1)}$, $L^{-1} \in \mathcal{A} \otimes_{\mathcal{O}} \underline{\mathcal{D}} \otimes_{\mathcal{O}} \mathcal{B}^{\otimes(-1)}$ and therefore $uL^{-1}v^t \in \underline{\mathcal{D}} \otimes_{\mathcal{O}} \Omega^{\otimes(-1)}$. The pairing (©.8) induces a pairing $\mathcal{E} \otimes \widetilde{\mathcal{E}} \to \mathcal{O}$ with the required properties. ■

©.2.3 Let us discuss the case $G = Sp(n)$. A $Sp(n)$-oper can be considered as a $GL(n)$-oper $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla)$ (see ©.2.1) with a horizontal symplective form on $\mathcal{E}$ such that $\mathcal{E}_i^{\perp} = \mathcal{E}_{n-i}$. We will construct an equivalence between the category of pairs $(\mathcal{A}, L)$ where $\mathcal{A}$ is an invertible sheaf on Y and L is a differential operator $\mathcal{A} \to \Omega \otimes \mathcal{A}^{\otimes(-1)}$ such that $L^t = L$ and the principle symbol σ of L does not vanish (in this case σ induces an isomorphism $\Omega^{\otimes n} \otimes \mathcal{A} \xrightarrow{\sim} \Omega \otimes \mathcal{A}^{\otimes(-1)}$ and therefore $\big(\mathcal{A} \otimes \Omega^{\otimes(n/2)}\big)^{\otimes 2} \xrightarrow{\sim} \Omega$, so $\mathcal{A} = \Omega^{\otimes(1/2)} \otimes \Omega^{\otimes(-n/2)}$ for some choice of $\Omega^{\otimes(1/2)}$.

If a pair $(\mathcal{A}, L)$ is given then the $GL(n)$-oper $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla)$ constructed in ©.2.1 is a $Sp(n)$-oper, the symplectic form on $\mathcal{E}$ being defined by (©.8). It is skew-symmetric because $L^t = L$ and therefore $\mathrm{res}\big(uL^{-1}v^t\big) = \mathrm{res}\big((vL^{-1}u^t)^t\big) = -\,\mathrm{res}\big(vL^{-1}u^t\big)$.

Conversely, let $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla)$ be a $Sp(n)$-oper and $L : \mathcal{A} \to \mathcal{B}$ the differential operator constructed from $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla)$ in ©.2.1. According to (©.5) $\mathcal{A} = \mathcal{E}_1^{\otimes(-1)}$ and $\mathcal{B} = \Omega \otimes (\mathcal{E}/\mathcal{E}_{n-1})^{\otimes(-1)}$. The pairing $\mathcal{E}/\mathcal{E}_{n-1} \otimes \mathcal{E}_1 \to \mathcal{O}$ induced by $\mathcal{E} \otimes \mathcal{E} \to \mathcal{O}$ induces an isomorphism $(\mathcal{E}/\mathcal{E}_{n-1})^* \xrightarrow{\sim} \mathcal{E}_1$ and therefore $\mathcal{B} \xrightarrow{\sim} \Omega \otimes \mathcal{A}^{\otimes(-1)}$. So we can consider L as an operator $\mathcal{A} \to \Omega \otimes \mathcal{A}^{\otimes(-1)}$. To show that $L^t = L$ notice that according to Proposition ©.2.2 the element of $\mathrm{Op}_n\big(\mathcal{A}, \Omega \otimes \mathcal{A}^{\otimes(-1)}\big)$ corresponding to $-L^t \in \mathrm{Diff}_n\big(\mathcal{A}, \Omega \otimes \mathcal{A}^{\otimes(-1)}\big)$ is the $GL(n)$-oper $(\mathcal{E}^*, \{(\mathcal{E}^*)_i\}, \nabla)$ with the isomorphisms $\varphi : (\mathcal{E}^*)_1 \xrightarrow{\sim} \mathcal{A}^{\otimes(-1)}$

and $\psi : \mathcal{E}^*/(\mathcal{E}^*)_{n-1} \xrightarrow{\sim} \mathcal{A}$ defined as the compositions $(\mathcal{E}^*)_1 \xrightarrow{\sim} (\mathcal{E}/\mathcal{E}_{n-1})^* \xrightarrow{\sim} \mathcal{E}_1 \xrightarrow{\sim} \mathcal{A}^{\otimes(-1)}$ and $\mathcal{E}^*/(\mathcal{E}^*)_{n-1} \xrightarrow{\sim} (\mathcal{E}_1)^* \xrightarrow{\sim} \mathcal{A}$. Since $-L^t$ corresponds to $(\mathcal{E}^*, \{(\mathcal{E})_i^*\}, \nabla, \varphi, \psi)$ the operator L^t corresponds to $(\mathcal{E}^*, \{(\mathcal{E})_i^*\}, \nabla, -\varphi, \psi)$. The pairing $\mathcal{E} \otimes \mathcal{E} \to \mathcal{O}$ induces an isomorphism $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla) \xrightarrow{\sim} (\mathcal{E}^*, \{(\mathcal{E}^*)_i\}, \nabla)$ such that diagrams

$$\begin{array}{ccc} \mathcal{E}_1 & \xrightarrow{\sim} & (\mathcal{E}^*)_1 \\ & {\scriptstyle\sim}\searrow \quad \swarrow{\scriptstyle\sim -\varphi} & \\ & \mathcal{A}^{\otimes(-1)} & \end{array} \qquad \begin{array}{ccc} \mathcal{E}/\mathcal{E}_{n-1} & \xrightarrow{\sim} & \mathcal{E}^*/(\mathcal{E}^*)_{n-1} \\ & {\scriptstyle\sim}\searrow \quad \swarrow{\scriptstyle\sim\psi} & \\ & \mathcal{A} & \end{array}$$

are commutative. So $L^t = L$.

©.2.4 According to ©.1.4 $sp(n)$-opers may be considered as $Sp(n)$-opers modulo twisting by Z-torsors where Z is the center of $Sp(n)$. According to ©.2.3 an $Sp(n)$-oper is the same as a pair $(\mathcal{M}, L)$ where $\mathcal{M}$ is a square root of Ω and L is a differential operator $\mathcal{M} \otimes \Omega^{\otimes(-n/2)} \to \mathcal{M} \otimes \Omega^{\otimes(n/2)}$ of order n such that $L^t = L$ and the principal symbol of L equals 1. Denote by $S(\mathcal{M})$ the set of all such L. If $\mathcal{M} \otimes \mathcal{A}$ is another square root of Ω then $\mathcal{A}$ is a square root of $\mathcal{O}$ and the canonical connection on $\mathcal{A}$ induces a bijection $S(\mathcal{M}) \xrightarrow{\sim} S(\mathcal{M} \otimes \mathcal{A})$. Replacing $\mathcal{M}$ by $\mathcal{M} \otimes \mathcal{A}$ and $L \in S(\mathcal{M})$ by its image in $S(\mathcal{M} \otimes \mathcal{A})$ corresponds to twisting $Sp(n)$-opers by Z-torsors.

So the set of $sp(n)$-opers can be identified with the set of differential operators $L : \Omega^{\otimes(1/2)} \otimes \Omega^{\otimes(-n/2)} \to \Omega^{\otimes(1/2)} \otimes \Omega^{\otimes(n/2)}$ of order n such that $L^t = L$ and the principal symbol of L equals 1 $\big($the latter set does not depend on the choice of $\Omega^{\otimes(1/2)}\big)$.

©.2.5 Let $n \in \mathbb{N}$ be odd. By an $O(n)$-oper we mean a $GL(n)$-oper $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla)$ (see ©.2.1) with a horizontal non-degenerate symmetric bilinear form B on $\mathcal{E}$ such that $\mathcal{E}_i^\perp = \mathcal{E}_{n-i}$. An $SO(n)$-oper in the sense of ©.1.2 can be considered as an $O(n)$-oper $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla, B)$ with an isomorphism $\alpha : \mathcal{O} \xrightarrow{\sim} \det \mathcal{E}$ compatible with the bilinear form on $\det \mathcal{E}$ induced by B. If L is a differential operator $\Omega^{\otimes(1-n)/2} \to \Omega^{\otimes(1+n)/2}$ of order n such that $L^t = -L$ and the principal symbol of L equals 1 then we construct an $SO(n)$-oper as follows. Take the $GL(n)$-oper $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla)$ corresponding to L according to ©.2.1 and define the bilinear form B on $\mathcal{E}$ by (©.8). Thus we obtain an $O(n)$-oper. The canonical isomorphisms $\mathcal{E}_i/\mathcal{E}_{i-1} \xrightarrow{\sim} \Omega^{\otimes(n+1-2i)/2}$ $\big($cf. (©.5)$\big)$ induce an isomorphism $\alpha : \mathcal{O} \xrightarrow{\sim} \det \mathcal{E}$. To show that it is compatible with the bilinear form on $\det \mathcal{E}$ induced by B we may assume that there

is a regular function z on Y such that dz has no zeros. Then Ω is trivial and $\mathcal{E}$ can be identified with the sheaf of differential operators on Y of order $\leq n-1$. So there is a base $e_i = \partial_z^i \in H^0(Y, \mathcal{E})$, $0 \leq i \leq n-1$. The isomorphism $\alpha : \mathcal{O} \xrightarrow{\sim} \det \mathcal{E}$ is defined by $1 \mapsto e_0 \wedge \ldots \wedge e_{n-1}$ and we have to show that the matrix $B_{ij} = B(e_i, e_j)$ has determinant 1. Indeed, $B_{ij} = 0$ if $i + j < n-1$ and $B_{ij} = (-1)^i$ if $i + j = n-1$, so $\det(B_{ij}) = 1$.

Conversely, if $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla, B, \alpha)$ is an $SO(n)$-oper then we have canonical isomorphisms $\mathcal{E}_{k+1}/\mathcal{E}_k \xrightarrow{\sim} \det \mathcal{E} \xrightarrow{\sim} \mathcal{O}$ where $k = (n-1)/2$. By virtue of (©.5) we have canonical isomorphisms $\mathcal{E}_1 \xrightarrow{\sim} \Omega^{\otimes(n-1)/2}$, $\mathcal{E}/\mathcal{E}_{n-1} \xrightarrow{\sim} \Omega^{\otimes(1-n)/2}$. So $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla)$ defines an operator $L \in \mathrm{Diff}_n\left(\Omega^{\otimes(1-n)/2}, \Omega^{\otimes(1+n)/2}\right)$ whose principal symbol equals 1. The argument used in ©.2.3 shows that $L^t = -L$.

So for an odd n we have constructed a bijection between the set of $SO(n)$-opers and the set of differential operators $L : \Omega^{\otimes(1-n)/2} \to \Omega^{\otimes(1+n)/2}$ of order n such that $L^t = -L$ and the principal symbol of L equals 1.

©.2.6 Since $sl(2) = sp(2) = o(3)$ one can interpret $sl(2)$-opers in two different ways.

1) According to ©.2.4 $sp(2)$-opers on Y bijectively correspond to Sturm-Liouville operators $L : \Omega^{\otimes(1/2)} \otimes \Omega^{\otimes(-1)} \to \Omega^{\otimes(1/2)} \otimes \Omega$ for a fixed choice of $\Omega^{\otimes(1/2)}$. By a Sturm-Liouville operator we mean a differential operator L of order 2 such that $L^t = L$ and the principal symbol of L equals 1. The set of Sturm-Liouville operators $\Omega^{\otimes(1/2)} \otimes \Omega^{\otimes(-1)} \to \Omega^{\otimes(1/2)} \otimes \Omega$ "does not depend" on the choice of $\Omega^{\otimes(1/2)}$ (for different choices of $\Omega^{\otimes(1/2)}$ there is a canonical bijection between the corresponding sets of Sturm-Liouville operators). Sturm-Liouville operators are also called *projective connections* on Y (we remind the origin of this term in ©.2.7).

2) According to ©.2.5 $o(3)$-opers (i.e., $SO(3)$-opers) bijectively correspond to differential operators $\widetilde{L} : \Omega^{\otimes(-1)} \to \Omega^{\otimes(2)}$ of order 3 such that $\widetilde{L}^t = -L$ and the principal symbol of $\widetilde{L}$ equals 1.

We are going to describe the relation between the operators L and $\widetilde{L}$ corresponding to the same $sl(2)$-oper. We will also give a third interpretation of $sl(2)$-opers.

Let us introduce some notation. If $\mathcal{L}$ and $\mathcal{M}$ are $\mathcal{O}$-modules set

$$\mathcal{L}\boxtimes\mathcal{M} = p_1^*\mathcal{L}\otimes p_2^*\mathcal{M} \tag{©.9}$$

where p_1 and p_2 are the projections $Y\times Y\to Y$. If $a,b\in\mathbb{Z}$, $a\geq b$, then set

$$\mathcal{P}_{a,b}(\mathcal{L},\mathcal{M}) = (\mathcal{L}\boxtimes\mathcal{M})(a\Delta)/(\mathcal{L}\boxtimes\mathcal{M})(b\Delta) \tag{©.10}$$

where $\Delta\subset Y\times Y$ is the diagonal.

If $\mathcal{E}$ and $\mathcal{F}$ are locally free $\mathcal{O}$-modules denote by $\mathrm{Diff}_{\leq n}(\mathcal{E},\mathcal{F})$ the space of differential operators $\mathcal{E}\to\mathcal{F}$ of order $\leq n$. There is a well known canonical isomorphism

$$\mathrm{Diff}_{\leq n}(\mathcal{E},\mathcal{F})\xrightarrow{\sim}H^0\big(Y\times Y,\mathcal{P}_{n+1,0}(\Omega\otimes\mathcal{E}^*,\mathcal{F})\big) \tag{©.11}$$

characterized by the following property: if $K(x,y)$ is a section of $\mathcal{P}_{n+1,0}\big(\Omega\otimes\mathcal{E}^*,\mathcal{F}\big)$ then the corresponding $L\in\mathrm{Diff}_{\leq n}(\mathcal{E},\mathcal{F})$ is defined by

$$(Ls)(y) = n!\operatorname*{res}_{x=y}\big(K(x,y)s(x)\big) \tag{©.12}$$

(i.e., L is the "integral operator" corresponding to the "kernel" $n!K$). Replacing $K(x,y)$ by $K(y,x)$ is equivalent to replacing $L:\mathcal{E}\to\mathcal{F}$ by $-L^t:\Omega\otimes\mathcal{F}^*\to\Omega\otimes\mathcal{E}^*$. Notice that due to the factor $n!$ in (©.10) the diagram

$$\begin{array}{ccc}\mathrm{Diff}_{\leq n}(\mathcal{E},\mathcal{F}) & \xrightarrow{\sim} & H^0\big(Y\times Y,\mathcal{P}_{n+1,0}(\Omega\otimes\mathcal{E}^*,\mathcal{F})\big)\\ {\scriptstyle\sigma}\big\downarrow & & \big\downarrow\\ H^0\big(Y,\mathcal{E}^*\otimes\Omega^{\otimes(-n)}\otimes\mathcal{F}\big) & \xleftarrow[\sim]{\varphi} & H^0\big(Y\times Y,\mathcal{P}_{n+1,n}(\Omega\otimes\mathcal{E}^*,\mathcal{F})\big)\end{array} \tag{©.13}$$

is commutative where σ associates to $L\in\mathrm{Diff}_{\leq n}(\mathcal{E},\mathcal{F})$ its principal symbol and φ is induced by the isomorphism $\mathcal{O}_{Y\times Y}\big((n+1)\Delta\big)/\mathcal{O}_{Y\times Y}(n\Delta)\xrightarrow{\sim}\Omega^{\otimes(-n-1)}$ defined by $(x-y)^{-n-1}f(x,y)\mapsto f(x,x)(dx)^{-n-1}$ (here we consider $\mathcal{O}_{Y\times Y}\big((n+1)\Delta\big)/\mathcal{O}_{Y\times Y}(n\Delta)$ as a sheaf on $\Delta=Y$).

Now we can describe the relation between the Sturm-Liouville operator L and the operator $\widetilde{L}$ of order 3 corresponding to the same $sl(2)$-oper. As explained above L corresponds to a skew-symmetric section of $\mathcal{P}_{3,0}\big(\Omega^{\otimes(3/2)},\Omega^{\otimes(3/2)}\big)$, which can be uniquely lifted to a skew-symmetric $K\in H^0\big(Y\times Y,\mathcal{P}_{3,-1}(\Omega^{\otimes(3/2)},\Omega^{\otimes(3/2)})\big)$. On the other hand $\widetilde{L}$ corresponds to a symmetric $\widetilde{K}\in H^0\big(Y\times Y,\mathcal{P}_{4,0}(\Omega^{\otimes 2},\Omega^{\otimes 2})\big)$. Both K and $\widetilde{K}$ have standard principal parts: $K(x,y)=(x-y)^{-3}(dx)^{3/2}(dy)^{3/2}+\ldots$, $\widetilde{K}(x,y)=(x-y)^{-4}(dx)^2(dy)^2+\ldots$.

Proposition. $\widetilde{K} = K^{4/3}$.

Proof. Suppose that in terms of a local parameter z the operator L is written as $\partial_z^2 + u$ $\big($i.e., $L\big(f(dz)^{-1/2}\big) = \big(\partial_z^2 f + uf\big)(dz)^{3/2}\big)$. The corresponding $sl(2)$-oper is defined by the connection

$$\partial_z + \begin{pmatrix} 0 & -u \\ 1 & 0 \end{pmatrix} \tag{©.14}$$

By $o(3)$ we understand the Lie algebra of the automorphism group of the quadratic form corresponding to the matrix

$$\begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

The image of (©.14) under a suitable isomorphism $sl(2) \xrightarrow{\sim} o(3)$ equals to

$$\partial_z + \begin{pmatrix} 0 & -2u & 0 \\ 1 & 0 & -2u \\ 0 & 1 & 0 \end{pmatrix} \tag{©.15}$$

The $o(3)$-oper defined by (©.15) corresponds to $\widetilde{L} = \partial_z^3 + 2u \cdot \partial_z + 2\partial_z \cdot u$. Recall that $2K$ and $6\widetilde{K}$ are the "kernels" of L and $\widetilde{L}$. So

$$K = \left\{ \frac{1}{(z_1 - z_2)^3} + \frac{u(z_1) + u(z_2)}{4(z_1 - z_2)} + \dots \right\} (dz_1)^{3/2} (dz_2)^{3/2} \tag{©.16}$$

$$\widetilde{K} = \left\{ \frac{1}{(z_1 - z_2)^4} + \frac{u(z_1) + u(z_2)}{3(z_1 - z_2)^2} + \dots \right\} (dz_1)^2 (dz_2)^2 \tag{©.17}$$

Therefore $\widetilde{K} = K^{4/3}$. ■

In ©.2.7 we will sketch a non-computational proof of the formula $\widetilde{K} = K^{4/3}$.

Now we can give the third interpretation of $sl(2)$-opers: they are in bijective correspondence with symmetric $\mathcal{K} \in H^0\big(Y \times Y, \mathcal{P}_{2,-2}(\Omega, \Omega)\big)$ with the standard principle part $\mathcal{K}(x, y) = (x - y)^{-2} dx\, dy + \dots$. The bijection is defined by

$$\mathcal{K} = \widetilde{K}^{1/2} = K^{2/3} \tag{©.18}$$

where $2K$ is the "kernel" of the Sturm-Liouville operator $L : \Omega^{\otimes(-1/2)} \to \Omega^{\otimes(3/2)}$ corresponding to the $sl(2)$-oper and $6\widetilde{K}$ is the "kernel" of the corresponding differential operator $\widetilde{L} : \Omega^{\otimes(-1)} \to \Omega^{\otimes 2}$ of order 3.

©.2.7 In this section we remind various facts about $sl(2)$-opers while in ©.2.8 and ©.2.9 we give an interpretation of $SL(n)$-opers and $SO(2k)$-opers. The reader can skip ©.2.7–©.2.9 and pass directly to ©§3.

In the previous sections we considered *algebraic* opers. If Y is an analytic curve over $\mathbb{C}$ and the words "bundle" and "connection" in Definition ©.1.2 are understood in the analytic sense then one obtains the notions of *analytic G-oper* and *analytic $\mathfrak{g}$-oper*. An analytic $sl(2)$-oper on Y is the same as a *locally-projective structure* on Y. Indeed, an $sl(2)$-oper (i.e., a $PGL(2)$-oper) can be considered as a quadruple (E, π, s, ∇) where $\pi : E \to Y$ is a locally trivial bundle with $\mathbb{P}^1$ as a fiber, ∇ is a connection on it and $s : Y \to E$ is a section such that the covariant differential of s does not vanish. In the analytic situation this is the same as locally-projective structure. This explains the term "projective connection" used as a synonym for "Sturm-Liouville operator" or "$sl(2)$-oper." In terms of the Sturm-Liouville operator L "good" local coordinates on Y are defined as ψ_1/ψ_2 where ψ_1 and ψ_2 are local solutions to the equation $L\psi = 0$; two good local coordinates z and ζ are related by a projective transformation $z = (a\zeta + b)/(c\zeta + d)$.

In the analytic situation the relation between the differential operators $L : \Omega^{\otimes(-1/2)} \to \Omega^{\otimes(3/2)}$ and $\widetilde{L} : \Omega^{\otimes(-1)} \to \Omega^{\otimes 2}$ can be expressed in the following way: the sheaf $\operatorname{Ker} \widetilde{L}$ is generated by the products $\psi_1\psi_2$ where ψ_1 and ψ_2 are local sections of $\operatorname{Ker} L$.

Notice that $\Omega^{\otimes(-1)} = \Theta$ is the tangent sheaf and therefore has a Lie algebra structure. In fact $\operatorname{Ker} \widetilde{L} \subset \Theta$ is a Lie subalgebra which is nothing but the sheaf of vector fields preserving L (i.e., preserving the locally-projective structure corresponding to L). This is an immediate consequence of the following expression for $\widetilde{L}$ in terms of L which holds in the algebraic situation as well as in the analytic one:

$$\widetilde{L}(v) = 2[v, L] \qquad (©.19)$$

Here $v \in \Theta = \Omega^{\otimes(-1)}$ (i.e., v is a local section of Θ) and $[v, L]$ is the Lie derivative of L with respect to v $\big(\Theta$ acts on $\Omega^{\otimes(-1)}$, $\Omega^{\otimes(3/2)}$, and therefore on the sheaf of differential operators $\Omega^{\otimes(-1/2)} \to \Omega^{\otimes(3/2)}$; in fact the differential operator $[v, L]$ has order 0, i.e., $[v, L] \in \Omega^{\otimes 2}\big)$.

Remark. Proposition ©.2.6 can be proved without computation. Indeed, the formula $L \mapsto \widetilde{K} - K^{4/3}$ defines a mapping $\varphi : \{\textit{projective connections}\} \longrightarrow \{\textit{quadratic differentials}\}$. To show that for any L the quadratic differential $\omega = \varphi(L)$ vanishes at any $y \in Y$ introduce a "good" local parameter ζ at y, i.e., $\zeta = \psi_1/\psi_2$ where ψ_1 and ψ_2 are local solutions to $L\psi = 0$ such that $\psi_1(y) = 0$. Then $L\big(f \cdot (d\zeta)^{-1/2}\big) = f''(d\zeta)^{3/2}$ and L is invariant under $\zeta \mapsto a\zeta$. So ω is also invariant and therefore $\omega(y) = 0$. This proof is in fact purely algebraic (one can consider formal solutions to $L\psi = 0$ instead of analytic ones).

©.2.8 An $SL(n)$-oper can be considered as a $GL(n)$-oper $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla)$ with a fixed horizontal isomorphism

$$\mathcal{O} \xrightarrow{\sim} \det \mathcal{E} \tag{©.20}$$

As explained in ©.2.1 $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla)$ defines a differential operator $L : \mathcal{A} \to \mathcal{A} \otimes \Omega^{\otimes n}$ or order n with principal symbol 1 where $\mathcal{A} = \mathcal{E}_1^{\otimes(-1)}$. The isomorphisms (©.5) and (©.20) induce an isomorphism

$$\mathcal{A}^{\otimes n} \otimes \Omega^{\otimes n(n-1)/2} \xrightarrow{\sim} \mathcal{O} \tag{©.21}$$

L^t is an operator $\mathcal{A} \otimes \mathcal{M} \to \mathcal{A} \otimes \Omega^{\otimes n} \otimes \mathcal{M}$ where $\mathcal{M} = \Omega^{\otimes(1-n)} \otimes \mathcal{A}^{\otimes -2}$. The isomorphism (©.21) induces an isomorphism $\mathcal{M}^{\otimes n} \xrightarrow{\sim} \mathcal{O}$ and therefore a connection on $\mathcal{M}$. So L^t induces a differential operator $\mathcal{A} \to \mathcal{A} \otimes \Omega^{\otimes n}$ which we also denote by L^t.

Proposition. *The order of $L + (-1)^{n+1} L^t$ is less than $n-1$ (i.e., the subprincipal symbol of L equals 0). Thus one obtains an equivalence between the category of $SL(n)$-opers and the category of pairs $(\mathcal{A}, L)$ where $\mathcal{A}$ is an invertible sheaf on Y with an isomorphism (©.21) and $L : \mathcal{A} \to \mathcal{A} \otimes \Omega^{\otimes n}$ is a differential operator of order n with principal symbol 1 and subprincipal symbol 0.*

The proof is left to the reader. It is based on the following local observation. Assume that there is a regular function z on Y such that dz has no zeros. Let $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla)$ be the $GL(n)$-oper corresponding to the operator $L : \mathcal{O} \to \mathcal{O}$, $L = \partial_z^n + \sum_{i=0}^{n-1} f_i \partial_z^i$. Identify $\mathcal{E}$ with the sheaf of differential operators on Y of order $\leq n-1$. Then the matrix q of the connection ∇ in the basis $e_i = \partial_z^i \in H^0(Y, \mathcal{E})$, $0 \leq i \leq n-1$, equals

$$-\begin{pmatrix} 0 \dots & 0 & f_0 \\ \dots & \dots & \dots \\ 0 \dots & 0 & f_{n-1} \end{pmatrix}$$

and so $Trq = -f_{n-1}$.

Clearly there is a natural bijection between the set of $sl(n)$-opers and the set of differential operators $L : \Omega^{\otimes(1-n)/2} \to \Omega^{\otimes(1+n)/2}$ of order n with principal symbol 1 and subprincipal symbol 0 (the latter set does not depend on the choice of $\Omega^{\otimes(1/2)}$).

©.2.9 Let us describe $SO(n)$-opers and $so(n)$-opers for $n = 2k$, $k \in \mathbb{Z}$, $k \geq 2$. By an $SO(n)$ vector bundle we mean a locally free $\mathcal{O}$-module $\mathcal{F}$ of rank n with a non-degenerate symmetric bilinear form B on it and an isomorphism $\alpha : \mathcal{O} \xrightarrow{\sim} \det \mathcal{F}$ compatible with the bilinear form $\overline{B}$ on $\det \mathcal{F}$ induced by B $\big($if Y were a curve over a field which is not algebraically closed then it would probably be more reasonable to require α to be compatible with $(-1)^k \overline{B}$ because the determinant of the hyperbolic symmetric bilinear form on $\mathbb{Z}^{2k}$ equals $(-1)^k\big)$. An $SO(n)$-oper is an $SO(n)$ vector bundle $\mathcal{F}$ with a filtration $\mathcal{F} = \mathcal{F}_{n-1} \supset \mathcal{F}_{n-2} \supset \ldots \supset \mathcal{F}_1 \supset \mathcal{F}_0 = 0$ and a connection $\nabla : \mathcal{F} \to \mathcal{F} \otimes \Omega$ compatible with B such that

1) the sheaves $\mathcal{F}_i/\mathcal{F}_{i-1}$ are invertible for $1 \leq i \leq n-1$, $i \neq k$ while $\mathcal{F}_k/\mathcal{F}_{k-1}$ is locally free of rank 2;

2) $\mathcal{F}_i^{\perp} = \mathcal{F}_{n-1-i}$;

3) $\nabla(\mathcal{F}_i) \subset \mathcal{F}_{i+1} \otimes \Omega$ for $1 \leq i \leq n-2$;

4) ∇ induces an isomorphism $\mathcal{F}_i/\mathcal{F}_{i-1} \xrightarrow{\sim} \big(\mathcal{F}_{i+1}/\mathcal{F}_i\big) \otimes \Omega$ for $1 \leq i \leq n-2$, $i \neq k-1$, k;

5) the composition of the morphisms

$$\mathcal{F}_{k-1}/\mathcal{F}_{k-2} \to \big(\mathcal{F}_k/\mathcal{F}_{k-1}\big) \otimes \Omega \to \big(\mathcal{F}_{k+1}/\mathcal{F}_k\big) \otimes \Omega^{\otimes 2}$$

induced by ∇ is an isomorphism.

Let $L : \Omega^{\otimes(1-k)} \to \Omega^{\otimes k}$ be a differential operator of order $2k-1$ such that $L^t = -L$ and the principal symbol of L equals 1. Starting from L and a section $f \in H^0\big(Y, \Omega^{\otimes k}\big)$ one can construct an $SO(n)$-oper in the following way. Let $(\mathcal{E}, \{\mathcal{E}_i\}, \nabla_{\mathcal{E}})$ be the $SO(n-1)$-oper corresponding to L (see ©.2.5). According to (©.5) we have canonical isomorphisms $\mathcal{E}_i/\mathcal{E}_{i-1} \xrightarrow{\sim} \Omega^{\otimes(k-i)}$ and in particular $\mathcal{E}_1 \xrightarrow{\sim} \Omega^{\otimes(k-1)}$. Denote by $\mathcal{F}$ the orthogonal direct sum $\mathcal{E} \oplus \mathcal{O}$, the bilinear form on $\mathcal{O}$ being standard. There is a unique connection $\nabla : \mathcal{F} \to \mathcal{F} \otimes \Omega$ compatible with the bilinear form on $\mathcal{F}$ such that the composition $\mathcal{E} \hookrightarrow \mathcal{F} \xrightarrow{\nabla} \mathcal{F} \otimes \Omega \to \mathcal{E} \otimes \Omega$ is equal to $\nabla_{\mathcal{E}}$ and $\nabla(1) = f$ where $1 \in H^0(Y, \mathcal{O}) \subset H^0(Y, \mathcal{F})$ and $f \in H^0\big(Y, \Omega^{\otimes k}\big) = H^0(Y, \mathcal{E}_1 \otimes \Omega) \subset H^0(Y, \mathcal{F} \otimes \Omega)$. Set $\mathcal{F}_i = \mathcal{E}_i$ for $i \leq k-1$, $\mathcal{F}_i = \mathcal{E}_i \oplus \mathcal{O}$ for $i \leq k$. The isomorphism $\mathcal{O} \xrightarrow{\sim} \det \mathcal{E}$ induces an isomorphism $\mathcal{O} \xrightarrow{\sim} \det \mathcal{F}$. So we have obtained a structure of $SO(n)$-oper on $\mathcal{F}$. This $SO(n)$-oper will be denoted by $\mathcal{F}(L, f)$. If $\mathcal{M}$ is a square root of $\mathcal{O}$ $\big($i.e., an invertible sheaf with a fixed isomorphism $\mathcal{O} \xrightarrow{\sim} \mathcal{M}^{\otimes 2}\big)$ then $\mathcal{F}(L, f) \otimes \mathcal{M}$ is also an $SO(n)$-oper.

Proposition. *Every $SO(n)$-oper is isomorphic to $\mathcal{F}(L,f)\otimes\mathcal{M}$ for some L, f and $\mathcal{M}$. $\mathcal{F}(L_1,f_1)\otimes\mathcal{M}_1$ and $\mathcal{F}(L_2,f_2)\otimes\mathcal{M}_2$ are isomorphic if and only if $L_1=L_2$, $f_1=f_2$ and there exists an isomorphism $\varphi:\mathcal{M}_1\xrightarrow{\sim}\mathcal{M}_2$ such that the diagram*

$$\begin{array}{ccc} & \mathcal{O} & \\ {\scriptstyle\sim}\swarrow & & \searrow{\scriptstyle\sim} \\ \mathcal{M}_1\otimes\mathcal{M}_1 & \xrightarrow[\sim]{\varphi\otimes\varphi} & \mathcal{M}_2\otimes\mathcal{M}_2 \end{array} \qquad (\text{©}.22)$$

is commutative. Moreover, isomorphisms $\mathcal{F}(L,f)\otimes\mathcal{M}_1\xrightarrow{\sim}\mathcal{F}(L,f)\otimes\mathcal{M}_2$ bijectively correspond to isomorphisms $\varphi:\mathcal{M}_1\xrightarrow{\sim}\mathcal{M}_2$ such that (©.22) is commutative.

Corollary. *There is a natural bijection between the set of $so(n)$-opers and the set of pairs (L,f) where $f\in H^0\big(Y,\Omega^{\otimes k}\big)$ and $L:\Omega^{\otimes(1-k)}\to\Omega^{\otimes k}$ is a differential operator of order $2k-1$ such that $L^t=-L$ and the principal symbol of L equals 1.*

Remark. As explained in [DS1], [DS2], it is natural to associate to (L,f) the expression $L+fd^{-1}f\in H^0\big(Y,\Omega^{\otimes k}\otimes_{\mathcal{O}}\underline{\mathcal{D}}\otimes_{\mathcal{O}}\Omega^{\otimes(k-1)}\big)$ where $\underline{\mathcal{D}}$ is the sheaf of pseudodifferential symbols and $d^{-1}\in H^0\big(Y,\underline{\mathcal{D}}\otimes_{\mathcal{O}}\Omega^{\otimes(-1)}\big)$ is inverse to the differential operator $d:\mathcal{O}\to\Omega$ considered as an element of $H^0\big(Y,\Omega\otimes_{\mathcal{O}}\mathcal{D}\big)$. One may say informally that $L+fd^{-1}f\in H^0\big(Y,\Omega^{\otimes k}\otimes_{\mathcal{O}}\underline{\mathcal{D}}\otimes_{\mathcal{O}}\mathcal{F}_1\big)$ annihilates $\mathcal{F}_1\subset\mathcal{F}$.

The proof of the Proposition is based on the following lemma.

Lemma. *Let $(\mathcal{F},\{\mathcal{F}_i\},\nabla)$ be an $SO(n)$-oper. Set $\mathcal{M}=\mathrm{Ker}\big(\nabla:\mathcal{F}_k/\mathcal{F}_{k-1}\to(\mathcal{F}_{k+1}/\mathcal{F}_k)\otimes\Omega\big)$. Then*

1) *the bilinear form on $\mathcal{F}_k/\mathcal{F}_{k-1}$ induces an isomorphism $\mathcal{M}\otimes\mathcal{M}\xrightarrow{\sim}\mathcal{O}$;*

2) *there is a unique way of lifting the inclusion $\mathcal{M}\hookrightarrow\mathcal{F}_k/\mathcal{F}_{k-1}$ to a morphism $s:\mathcal{M}\to\mathcal{F}_k$ such that $\nabla\big(s(\mathcal{M})\big)\subset\big(s(\mathcal{M})+\mathcal{F}_1\big)\otimes\Omega$.*

Using the lemma it is easy to prove the proposition by considering the orthogonal decomposition $\mathcal{F}=s(\mathcal{M})\oplus\mathcal{E}$.

Proof of the lemma. Since the composition of the morphisms $\mathcal{F}_{k-1}/\mathcal{F}_{k-2} \to (\mathcal{F}_k/\mathcal{F}_{k-1})\otimes\Omega \to (\mathcal{F}_{k+1}/\mathcal{F}_k)\otimes\Omega^{\otimes 2}$ induced by ∇ is an isomorphism $\mathcal{F}_k/\mathcal{F}_{k-1} = \mathcal{M}\oplus\mathcal{N}$ where $\mathcal{N}\otimes\Omega = \mathrm{Im}\big(\nabla : \mathcal{F}_{k-1}/\mathcal{F}_{k-2} \to (\mathcal{F}_k/\mathcal{F}_{k-1})\otimes\Omega\big)$. Since ∇ is compatible with the bilinear form $\mathcal{M}$ and $\mathcal{N}$ are orthogonal to each other. Since the bilinear form on $\mathcal{F}_k/\mathcal{F}_{k-1}$ is non-degenerate the statement 1) is clear. To prove the statement 2) we may assume that the curve Y is affine and there is an isomorphism $\mathcal{O}\xrightarrow{\sim}\mathcal{M}$ compatible with the bilinear form on $\mathcal{M}$ (otherwise replace $\mathcal{F}$ by $\mathcal{F}\otimes\mathcal{M}$). Choose $\sigma\in H^0(Y,\mathcal{F}_k)$ so that the image of σ in $H^0(Y,\mathcal{F}_k/\mathcal{F}_{k-1})$ equals $1\in H^0(Y,\mathcal{O}) = H^0(Y,\mathcal{M})\subset H^0(Y,\mathcal{F}_k/\mathcal{F}_{k-1})$. We must show that there is a unique $\tau\in H^0(Y,\mathcal{F}_{k-1})$ such that $\nabla(\sigma+\tau)\in\big(\mathcal{O}(\sigma+\tau)\oplus\mathcal{F}_1\big)\otimes\Omega$. Since $(\sigma+\tau,\sigma+\tau) = 1$ we have $\big(\nabla(\sigma+\tau),\sigma+\tau\big) = 0$ and therefore if $\nabla(\sigma+\tau)\in\big(\mathcal{O}(\sigma+\tau)\oplus\mathcal{F}_1\big)\otimes\Omega$ then $\nabla(\sigma+\tau)\in\mathcal{F}_1\otimes\Omega$. Now the uniqueness of τ is clear and the existence of τ means that

$$\nabla(\sigma)\in\big(\mathcal{F}_{k-1}\otimes\Omega\big) + \nabla\big(\mathcal{F}_{k-1}\big) \qquad (©.23)$$

Since the image of σ in $H^0(Y,\mathcal{F}_k)$ belongs to $\mathcal{M}$ we have $\nabla(\sigma)\in\mathcal{F}_k\otimes\Omega$ and since $\big(\nabla(\sigma),\sigma\big) = \frac{1}{2}d(\sigma,\sigma) = 0$ the image of $\nabla(\sigma)$ in $(\mathcal{F}_k/\mathcal{F}_{k-1})\otimes\Omega$ belongs to $\mathcal{N}\otimes\mathcal{M}$. This is equivalent to (©.23). ■

©§3. A description of the set of $\mathfrak{g}$-opers.

Denote by $\mathrm{Op}_{\mathfrak{g}}(Y)$ the set of isomorphism classes of $\mathfrak{g}$-opers on Y.

©.3.1 Let $\mathfrak{g}=\mathrm{sl}(2)$. In this case it is well known that $\mathrm{Op}_{\mathfrak{g}}(Y)$ is a principal homogeneous space over $H^0\big(Y,\Omega^{\otimes 2}\big)$. Let us recall why it is so. According to ©.2.6 $\mathrm{Op}_{\mathfrak{g}}(Y)$ can be interpreted as the set of Sturm-Liouville operators L on Y. So we obtain a free transitive action of $H^0\big(Y,\Omega^{\otimes 2}\big)$ on $\mathrm{Op}_{\mathfrak{g}}(Y)$ defined by

$$L\mapsto L+\omega,\quad \omega\in H^0\big(Y,\Omega^{\otimes 2}\big) \qquad (©.24)$$

Other interpretations of $\mathrm{Op}_{\mathfrak{g}}(Y)$ (see ©.2.6) give rise to actions of $H^0\big(Y,\Omega^{\otimes 2}\big)$ on $\mathrm{Op}_{\mathfrak{g}}(Y)$ which can be written in terms of the Sturm-Liouville operators as $L\mapsto L+\alpha\omega$ where $\alpha\in\mathbb{Q}$ depends on the interpretation.

Let us recall why $\mathrm{Op}_{\mathfrak{g}}(Y)\neq\emptyset$. Certainly there is an open covering $Y = \bigcup\limits_i U_i$ such that $\mathfrak{g}$-opers on $\mathcal{M}_i$ exist. So if $H^1\big(Y,\Omega^{\otimes 2}\big) = 0$ then $\mathrm{Op}_{\mathfrak{g}}(Y)\neq\phi$. If $H^1\big(Y,\Omega^{\otimes 2}\big)\neq 0$ then either Y is an elliptic curve or $Y=\mathbb{P}^1$. In the first case Ω is trivial and so the interpretation of $\mathfrak{g}$-opers as Sturm-Liouville operators shows that $\mathrm{Op}_{\mathfrak{g}}(Y)\neq\phi$. On $\mathbb{P}^1$ there is an obvious $SL(2)$-oper: take the trivial bundle $\mathcal{E}=\mathcal{O}^2$ with the trivial connection and define $\mathcal{E}_1\subset\mathcal{E}$ to be the tautological line subbundle. So $\mathrm{Op}_{\mathfrak{g}}(\mathbb{P}^1)\neq\phi$.

©.3.2 The action (©.24) can be described without using Sturm-Liouville operators. Let $(\mathcal{F},\nabla)$ be a $\mathfrak{g}$-oper in the sense of Definitions ©.1.2 and ©.1.3 where $\mathfrak{g}=sl(2)$. Then we can obtain a new $\mathfrak{g}$-oper $(\mathcal{F},\widetilde{\nabla})$ by setting $\widetilde{\nabla}=\nabla+\eta$, $\eta\in\mathrm{Hom}\big(\Theta,\mathfrak{g}^1_{\mathcal{F}}\big)=H^0\big(Y,\mathfrak{g}^1_{\mathcal{F}}\otimes\Omega\big)$ where $\mathfrak{g}^1_{\mathcal{F}}$ has the same meaning as in ©.1.1. We have the canonical isomorphism $\Theta\to\mathfrak{g}^{-1}_{\mathcal{F}}/\mathfrak{g}^0_{\mathcal{F}}$ induced by ∇ (see ©.1.2) and the pairing between $\mathfrak{g}^{-1}_{\mathcal{F}}/\mathfrak{g}^0_{\mathcal{F}}$ and $\mathfrak{g}^1_{\mathcal{F}}$ induced by the bilinear form. $Tr(XY)$ on $\mathfrak{g}$. So we can identify $\mathfrak{g}^1_{\mathcal{F}}$ with Ω. It is easy to see (cf. ©.1.4) that the action (©.24) corresponds to replacing ∇ by $\nabla-\omega$, $\omega\in H^0\big(Y,\Omega^{\otimes 2}\big)=H^0\big(Y,\mathfrak{g}^1_{\mathcal{F}}\otimes\Omega\big)$.

©.3.3 Let $B\subset SL(2)$ denote the subgroup of upper-triangular matrices and B_{ad} the image of B in $PGL(2)$. It follows from ©.3.1 and ©.3.2 that for any $sl(2)$-opers $(\mathcal{F},\nabla)$ and $(\widetilde{\mathcal{F}},\widetilde{\nabla})$ the B_{ad}-bundles $\mathcal{F}$ and $\widetilde{\mathcal{F}}$ are canonically isomorphic. This also follows from the construction of the $SL(2)$-oper corresponding to a Sturm-Liouville operator $\Omega^{\otimes(-1/2)}\to\Omega^{\otimes(-3/2)}$ (see ©.2.1 and ©.2.3 or ©.2.8). Indeed, the B-bundle in the construction corresponds to the $\mathcal{O}$-module $\mathcal{E}=\mathcal{D}_1\otimes_{\mathcal{O}}\Omega^{\otimes(1/2)}$ and the canonical exact sequence $0\to\Omega^{\otimes(1/2)}\to\mathcal{E}\to\Omega^{\otimes(-1/2)}\to 0$ (recall that $\mathcal{D}_1$ denotes the sheaf of differential operators of degree ≤ 1). This B-bundle depends on the choice of $\Omega^{\otimes(1/2)}$, but the corresponding B_{ad}-bundle is a quite canonical object.

Remark. In [De1] Deligne gave a nice description of this B_{ad}-bundle considered as a principal bundle: its fiber over a point $y\in Y$ is the set of $\mathbb{C}$-isomorphisms between $\mathrm{Spec}\,\mathcal{O}_y/m_y^3$ and $\mathrm{Spec}\,\mathcal{O}_\infty/m_\infty^3$ where $\mathcal{O}_y$ and $\mathcal{O}_\infty$ are the local rings of $y\in Y$ and $\infty\in\mathbb{P}^1$ while m_y and m_∞ are their maximal ideals (B_{ad} acts on this set because it acts on $\mathbb{P}^1$ preserving ∞). ■

©.3.4 Now let $\mathfrak{g}$ be an arbitrary semisimple Lie algebra with a fixed Borel subalgebra $\mathfrak{b}\subset\mathfrak{g}$ and a fixed Cartan subalgebra $\mathfrak{h}\subset\mathfrak{b}$. Let G_{ad} be the connected algebraic group with trivial center corresponding to $\mathfrak{g}$. Denote by H_{ad} and B_{ad} the Cartan and Borel subgroups of G_{ad} corresponding to $\mathfrak{h}$ and $\mathfrak{b}$. Fix an $sl(2)$-triple $\{h,x,y\}\subset\mathfrak{g}$ such that $h\in\mathfrak{h}$ and $x\in\mathfrak{b}$ is a principle nilpotent element. This means that $h=2\sum\limits_{\alpha\in\Gamma}\check{\omega}_\alpha$, $x=\sum\limits_{\alpha\in\Gamma}x_\alpha$, $y=\sum\limits_{\alpha\in\Gamma}y_\alpha$, where Γ is the set of simple roots, $\check{\omega}_\alpha$ are the fundamental coweights, $x_\alpha\in\mathfrak{g}^\alpha$, $y_\alpha\in\mathfrak{g}^{-\alpha}$, $x_\alpha\neq 0$, $y_\alpha\neq 0$. The y_α are uniquely determined by x_α because $[x,y]=h$ while the elements $x_\alpha\in\mathfrak{g}^\alpha$, $x_\alpha\neq 0$, can be arbitrary. However any two $sl(2)$-triples of the kind described above are conjugate by a unique element of H_{ad}.

Set $V = \operatorname{Ker} \operatorname{ad} x \subset \mathfrak{g}$. The grading $\mathfrak{g} = \underset{k}{\oplus}\mathfrak{g}_k$ (see ©.1.1) induces a grading $V = \underset{k}{\oplus}V_k$. It is well known that $V_k = 0$ if $k \leq 0$. The numbers k such that $V_k \neq 0$ are nothing but the exponents of $\mathfrak{g}$ and $\dim V$ is equal to the rank of $\mathfrak{g}$ (see [Ko1]). Notice that $V_1 \neq 0$ because $x \in V_1$. Of course V considered as a subspace of $\mathfrak{g}$ depends on $\{h, x, y\}$. But for any two choices of $\{h, x, y\}$ there is a canonical isomorphism between the corresponding spaces V. So we can identify all these spaces and consider the space $V(\mathfrak{g})$ which depends only on $\mathfrak{g}$ (of course $V(\mathfrak{g})$ is not a subspace of $\mathfrak{g}$). There is a canonical grading $V(\mathfrak{g}) = \underset{k}{\oplus}V_k(\mathfrak{g})$ and a canonical element $\mathcal{X} \in V_1(\mathfrak{g})$ corresponding to $x \in V_1$. Now set

$$V_k^{\mathfrak{g}}(Y) = V_{k-1}(\mathfrak{g}) \otimes H^0\big(Y, \Omega^{\otimes k}\big), \quad V^{\mathfrak{g}}(Y) = \underset{k}{\oplus}V_k^{\mathfrak{g}}(Y) \tag{©.25}$$

There is a canonical embedding $H^0\big(Y, \Omega^{\otimes 2}\big) \hookrightarrow V_2^{\mathfrak{g}}(Y) \hookrightarrow V^{\mathfrak{g}}(Y)$ defined by

$$\omega \mapsto -\omega \cdot \mathcal{X} \tag{©.26}$$

So the principal homogeneous space $\operatorname{Op}_{sl(2)}(Y)$ over $H^0\big(Y, \Omega^{\otimes 2}\big)$ induces a principal homogeneous space $\underline{\operatorname{Op}}_{\mathfrak{g}}(Y)$ over $V^{\mathfrak{g}}(Y)$. In other words $\underline{\operatorname{Op}}_{\mathfrak{g}}(Y)$ is the quotient of $\operatorname{Op}_{sl(2)}(Y) \times V^{\mathfrak{g}}(Y)$ modulo the equivalence relation

$$(L + \omega, \eta) \sim (L, \eta - \omega\mathcal{X}) \tag{©.27}$$

where $\eta \in V^{\mathfrak{g}}(Y)$, $\omega \in H^0\big(Y, \Omega^{\otimes 2}\big)$ and $L \in \operatorname{Op}_{sl(2)}(Y)$ is considered as a Sturm-Liouville operator. We will construct a canonical bijection $\underline{\operatorname{Op}}_{\mathfrak{g}}(Y) \xrightarrow{\sim} \operatorname{Op}_{\mathfrak{g}}(Y)$.

Remark. Since $\operatorname{Op}_{\mathfrak{g}_1 \times \mathfrak{g}_2}(Y) = \operatorname{Op}_{\mathfrak{g}_1}(Y) \times \operatorname{Op}_{\mathfrak{g}_2}(Y)$ it is enough to describe $\operatorname{Op}_{\mathfrak{g}}(Y)$ if $\mathfrak{g}$ is simple. In this case $V_1(\mathfrak{g})$ is generated by $\mathcal{X}$ and therefore $\underline{\operatorname{Op}}_{\mathfrak{g}}(Y) = \operatorname{Op}_{sl(2)}(Y) \times \big(\underset{k>2}{\oplus} V_k^{\mathfrak{g}}(Y)\big)$. ■

Let us construct a canonical mapping $\underline{\operatorname{Op}}_{\mathfrak{g}}(Y) \to \operatorname{Op}_{\mathfrak{g}}(Y)$. By definition, an $sl(2)$-oper L is a $PGL(2)$-oper $(\mathcal{F}_0, \nabla_0)$ where $\mathcal{F}_0$ is a torsor over the upper-triangular subgroup $B_0 \subset PGL(2)$ and ∇_0 is a connection on the corresponding $PGL(2)$-bundle satisfying a certain condition (see ©.1.2). The fixed $sl(2)$-triple $\{h, x, y\}$ defines a homomorphism $\rho : PGL(2) \to G_{\mathrm{ad}}$. Since $\rho(B_0) \subset B_{\mathrm{ad}}$ we can associate to $\mathcal{F}_0$ a B_{ad}-bundle $\mathcal{F}$ and ∇_0 induces a connection ∇ on the G_{ad}-bundle corresponding to $\mathcal{F}$. Clearly $(\mathcal{F}, \nabla)$ is a G_{ad}-oper, i.e., a $\mathfrak{g}$-oper. For $\eta \in \operatorname{Hom}(\Theta, \mathfrak{b}_{\mathcal{F}}) = H^0(Y, \mathfrak{b}_{\mathcal{F}} \otimes \Omega)$ the pair $(\mathcal{F}, \nabla + \eta)$ is still

a $\mathfrak{g}$-oper. Notice that $\mathfrak{b}_{\mathcal{F}} = \mathfrak{b}_{\mathcal{F}_0}$ where $\mathfrak{b}_{\mathcal{F}_0}$ corresponds to $\mathcal{F}_0$ and the representation of B_0 in $\mathfrak{b}$ defined as the composition of $\rho : B_0 \to B_{\mathrm{ad}}$ and the adjoint representation of B_{ad} in $\mathfrak{b}$. Since V is a B_0-invariant subspace of $\mathfrak{b}$ we have $\mathfrak{b}_{\mathcal{F}} = \mathfrak{b}_{\mathcal{F}_0} \supset V_{\mathcal{F}_0}$. An element $g \in B_0$ acts on V_k as multiplication by $\chi(g)^k$ where $\chi : B_0 \to \mathbb{G}_m$ maps $\left(\begin{smallmatrix} a & b \\ 0 & c \end{smallmatrix}\right)$ to a/c. Since the line bundle corresponding to $\mathcal{F}_0$ and χ is canonically isomorphic to Ω (see ©.1.4) we have $V_{\mathcal{F}_0} = \underset{k}{\oplus}\big(V_k \otimes \Omega^{\otimes k}\big)$. So $H^0\big(Y, \mathfrak{b}_{\mathcal{F}} \otimes \Omega\big) \subset H^0\big(Y, V_{\mathcal{F}_0} \otimes \Omega\big) = \underset{k}{\oplus}\big(V_k \otimes H^0(Y, \Omega^{\otimes(k+1)})\big) = V^{\mathfrak{g}}(Y)$. Therefore to an $sl(2)$-oper L and an $\eta \in V^{\mathfrak{g}}(Y)$ we can associate the $\mathfrak{g}$-oper $(\mathcal{F}, \nabla + \eta)$ where $(\mathcal{F}, \nabla)$ is constructed from L as explained above. It follows from ©.3.2 and (©.27) that the constructed mapping $\mathrm{Op}_{sl(2)}(Y) \times V^{\mathfrak{g}}(Y) \to \mathrm{Op}_{\mathfrak{g}}(Y)$ factors through $\underline{\mathrm{Op}}_{\mathfrak{g}}(Y)$. So we have obtained a canonical mapping

$$\underline{\mathrm{Op}}_{\mathfrak{g}}(Y) \to \mathrm{Op}_{\mathfrak{g}}(Y) \qquad (©.28)$$

Theorem. *The mapping (©.28) is bijective.*

Proof. Proposition ©.1.3 shows that the assertion is local. So it is enough to prove the following. Suppose that there is function $z \in H^0(Y, \mathcal{O}_Y)$ such that dz has no zeros and let q be a regular function on Y with values in $\mathfrak{g}^{-1} = \underset{k \geq -1}{\oplus} \mathfrak{g}_k$ such that for every simple root α the component $q_{-\alpha}$ of q has no zeros. Then there is a unique B_{ad}-valued regular function b on Y such the values of the function $\widetilde{q}$ defined by the relation $b^{-1}(\partial_z + q)b = \partial_z + \widetilde{q}$ belong to $y + V$ where y is the member of the $sl(2)$-triple $\{h, x, y\}$ and $V = \operatorname{Ker} \operatorname{ad} x$. This is easily deduced from the well known equalities $\mathfrak{g} = (\operatorname{Im} \operatorname{ad} y) \oplus V$ and $(\operatorname{Ker} \operatorname{ad} y) \cap \mathfrak{b} = 0$. ■

Though the bijection (©.28) is canonical we are not sure that it gives a reasonable description of $\mathrm{Op}_{\mathfrak{g}}(Y)$. So we will avoid using it when possible.

©.3.5 Let B_0, B_{ad} and $\rho : PGL(2) \to G_{\mathrm{ad}}$ denote the same objects as in ©.3.4. Denote by $\mathcal{F}_0^{\mathrm{can}}$ the canonical B_0-bundle on Y constructed in ©.3.3 and by $\mathcal{F}^{\mathrm{can}}$ the B-bundle corresponding to $\mathcal{F}_0^{\mathrm{can}}$ and the homomorphism $B_0 \to B$ induced by ρ. It follows from ©.3.3 and ©.3.4 that for any $\mathfrak{g}$-oper $(\mathcal{F}, \nabla)$ the B_{ad}-bundle $\mathcal{F}$ is canonically isomorphic to $\mathcal{F}^{\mathrm{can}}$.

©.3.6 Theorem ©.3.4 implies that for any finite subset $D \subset Y$ the natural mapping $\mathrm{Op}_{\mathfrak{g}}(Y) \to \mathrm{Op}_{\mathfrak{g}}(Y \backslash D)$ is injective. We will usually consider $\mathrm{Op}_{\mathfrak{g}}(Y)$ as a subset of $\mathrm{Op}_{\mathfrak{g}}(Y \backslash D)$.

Remark. The injectivity of the mapping $\mathrm{Op}_{\mathfrak{g}}(Y) \to \mathrm{Op}_{\mathfrak{g}}(Y \backslash D)$ also follows from the definition of $\mathfrak{g}$-oper and the following fact: if $\mathcal{E}_1$ and $\mathcal{E}_2$ are bundles with connections on Y and φ is a horizontal isomorphism between the restrictions of $\mathcal{E}_1$ and $\mathcal{E}_2$ to $Y \backslash D$ then φ is the restriction of an isomorphism $\mathcal{E}_1 \xrightarrow{\sim} \mathcal{E}_2$. However the argument based on Theorem ©.3.4 works in more general situations: for $\mathfrak{g}$-opers with singularities (see Proposition ©.4.2) and for $(\mathfrak{g}, h)$-opers (see ©.5.2).

©.3.7 The reader can skip this section and pass directly to ©§4.

As explained in ©.3.4 the space $V^{\mathfrak{g}}(Y)$ which appears in the definition of $\underline{\mathrm{Op}}_{\mathfrak{g}}(Y)$ is equal to $H^0\big(Y, V_{\mathcal{F}_0}\big)$ where $\mathcal{F}_0$ is the canonical torsor over the upper-triangular subgroup $B_0 \subset PGL(2)$ (see ©.3.3) and $V = \operatorname{Ker} \operatorname{ad} x \subset \mathfrak{g}$ is considered as a B_0 module via the embedding $B_0 \hookrightarrow B_{\mathrm{ad}}$ corresponding to $\{h, x, y\}$. Now let V' be any subspace of $\mathfrak{g}$ such that

$$[h, V'] \subset V', \quad [x, V'] \subset V', \quad \mathfrak{g} = (\operatorname{Im} \operatorname{ad} y) \oplus V', \quad x \in V' \tag{©.29}$$

Then one can construct an analog of $\underline{\mathrm{Op}}_{\mathfrak{g}}(Y)$ in which $V^{\mathfrak{g}}(Y) = H^0\big(Y, V_{\mathcal{F}_0}\big)$ is replaced by $H^0\big(Y, V'_{\mathcal{F}_0}\big)$ and an analog of the bijection (©.28). So it is natural to ask how to describe all subspaces $V' \subset \mathfrak{g}$ satisfying (©.29). If V' is such a subspace then we have the isomorphism $f : V' \xrightarrow{\sim} V$ defined as the composition $V' \xrightarrow{\sim} V/\operatorname{Im} \operatorname{ad} y \xrightarrow{\sim} V$. Therefore $\operatorname{ad} x : V' \to V'$ induces the operator $\xi : V \to V$, $\xi = f \circ (\operatorname{ad} x) \circ f^{-1}$. So we obtain a mapping from the set of all V' satisfying (©.29) to the set of all linear operators $\xi : V \to V$ such that

$$\xi(V_k) \subset V_{k+1}, \quad \xi(x) = 0 \tag{©.30}$$

It is easy to show that this mapping is a bijection. A glance into the list of exponents of simple Lie algebras shows that for simple Lie algebras $\mathfrak{g}$ of types B_n, C_n, D_{2k}, E_7, E_8, F_4, and G_2 the only possibility is $\xi = 0$, i.e., $V' = V$. For D_{2k+1} and E_6 there are some nontrivial choices of V' and for $\mathfrak{g} = sl(n)$ there are plenty subspaces V' satisfying (©.29). For instance one can take $V' = \mathbb{C}x \oplus \mathbb{C}e_{1n} \oplus \ldots \oplus \mathbb{C}e_{n-2,n}$ where e_{ij} is the matrix with 1 at the intersection of the i-th row and the j-th column and zeros elsewhere.

©§4. Singularities of $\mathfrak{g}$-opers.

Just as in ©§3 we denote by $\mathfrak{g}$ a semisimple Lie algebra and by $\mathrm{Op}_{\mathfrak{g}}(Y)$ the set of $\mathfrak{g}$-opers on Y. For every finite subscheme $D \subset Y$ we will introduce a certain subset $\mathrm{Op}_{\mathfrak{g},D}(Y) \subset \mathrm{Op}_{\mathfrak{g}}(Y\backslash D)$ such that $D \subset \Delta$ implies $\mathrm{Op}_{\mathfrak{g},D}(Y) \subset \mathrm{Op}_{\mathfrak{g},\Delta}(Y)$ and for every finite subset $S \subset Y$ $\mathrm{Op}_{\mathfrak{g}}(Y\backslash S)$ is the union of $\mathrm{Op}_{\mathfrak{g},D}(Y)$ for all D such that $\mathrm{Supp}\, D \subset S$. Here $\mathrm{Supp}\, D$ denotes D considered as a set.

©.4.1 In ©.3.4 we constructed a vector space $V^{\mathfrak{g}}(Y)$ (see (©.25)) and a canonical embedding $H^0(Y,\Omega^{\otimes 2}) \hookrightarrow V^{\mathfrak{g}}(Y)$ (see (©.26)). We defined $\underline{\mathrm{Op}}_{\mathfrak{g}}(Y)$ to be the principal homogeneous space over $V^{\mathfrak{g}}(Y)$ induced by $\mathrm{Op}_{sl(2)}(Y)$ via this embedding $\big($recall that $\mathrm{Op}_{sl(2)}(Y)$ is a principal homogenous space over $H^0(Y,\Omega^{\otimes 2})\big)$. Now for any finite subscheme $D \subset Y$ set

$$V^{\mathfrak{g}}_{D,k}(Y) = V_{k-1}(\mathfrak{g}) \otimes H^0\big(Y, \Omega^{\otimes k}(kD)\big), \quad V^{\mathfrak{g}}_D(Y) = \underset{k}{\oplus} V^{\mathfrak{g}}_{D,k}(Y) \tag{©.31}$$

where $V_{k-1}(\mathfrak{g})$ has the same meaning as in (©.25). Since $V^{\mathfrak{g}}(Y) \subset V^{\mathfrak{g}}_D(Y)$ the principal homogenous space $\underline{\mathrm{Op}}_{\mathfrak{g}}(Y)$ over $V^{\mathfrak{g}}(Y)$ induces a principle homogenous space over $V^{\mathfrak{g}}_D(Y)$ which will be denoted by $\underline{\mathrm{Op}}_{\mathfrak{g},D}(Y)$. If $D \subset \Delta$ then $\underline{\mathrm{Op}}_{\mathfrak{g},D}(Y) \subset \underline{\mathrm{Op}}_{\mathfrak{g},\Delta}(Y)$ and for any finite subset $S \subset Y$ $\underline{\mathrm{Op}}_{\mathfrak{g}}(Y\backslash S)$ is the union of $\underline{\mathrm{Op}}_{\mathfrak{g},D}(Y)$ for all D such that $\mathrm{Supp}\, D \subset S$.

©.4.2 We have a canonical bijection $\underline{\mathrm{Op}}_{\mathfrak{g}}(Y\backslash D) \xrightarrow{\sim} \mathrm{Op}_{\mathfrak{g}}(Y\backslash D)$ (see (©.28)). So we could define $\mathrm{Op}_{\mathfrak{g},D}(Y)$ as the image of $\underline{\mathrm{Op}}_{\mathfrak{g},D}(Y) \subset \underline{\mathrm{Op}}_{\mathfrak{g}}(Y\backslash D)$ in $\mathrm{Op}_{\mathfrak{g}}(Y\backslash D)$. But we prefer to give an equivalent "definition" of $\mathrm{Op}_{\mathfrak{g},D}(Y)$ which does not use the "suspicious" bijection (©.28).

Let G be the connected algebraic group with trivial center corresponding to $\mathfrak{g}$, B the Borel subgroup of G corresponding to $\mathfrak{b} \subset \mathfrak{g}$. In the following definition we use the notation of ©.1.1 and ©.1.2.

Definition. *A $\mathfrak{g}$-oper on Y with D-singularities is a B-bundle $\mathcal{F}$ on Y with a morphism $\nabla : \Theta(-D) \to \mathcal{E}^{-1}_{\mathcal{F}}$ such that the diagram*

$$\begin{array}{ccc} \Theta(-D) & \xrightarrow{\nabla} & \mathcal{E}^{-1}_{\mathcal{F}} \\ \big\downarrow & \swarrow & \\ \Theta & & \end{array} \tag{©.32}$$

is commutative and for every simple root α the composition $\Theta(-D) \xrightarrow{\nabla} \mathcal{E}^{-1}_{\mathcal{F}} \longrightarrow \mathcal{E}^{-1}_{\mathcal{F}}/\mathcal{E}_{\mathcal{F}} = \mathfrak{g}^{-1}_{\mathcal{F}}/\mathfrak{g}_{\mathcal{F}} \longrightarrow \mathfrak{g}^{-\alpha}_{\mathcal{F}}$ is an isomorphism.

Denote by $\mathrm{Op}_{\mathfrak{g},D}(Y)$ the set of isomorphism classes of $\mathfrak{g}$-opers on Y with D-singularities.

Proposition. *The natural mapping* $\mathrm{Op}_{\mathfrak{g},D}(Y) \to \mathrm{Op}_{\mathfrak{g}}(Y\backslash D)$ *is injective and its image is equal to the image of* $\underline{\mathrm{Op}}_{\mathfrak{g},D}(Y) \subset \underline{\mathrm{Op}}_{\mathfrak{g},D}(Y\backslash D)$ *in* $\mathrm{Op}_{\mathfrak{g}}(Y\backslash D)$.

Proof. Fix an $sl(2)$-triple $\{h, x, y\} \subset \mathfrak{g}$ such that $h \in \mathfrak{h}$ and $x \in \mathfrak{b}$ is a principle nilpotent element. Set $V = \operatorname{Ker} \operatorname{ad} x$. We may assume that there is function $z \in H^0(Y, \mathcal{O}_Y)$ such that dz has no zeros and that $D \subset Y$ is defined by the equation $f = 0$, $f \in H^0(Y, \mathcal{O}_Y)$. The main step of the proof is to show that if q is a regular function on Y with values in $\mathfrak{g}^{-1} = \underset{k\geq -1}{\oplus} \mathfrak{g}_k$ such that for every simple root α the component $q_{-\alpha}$ of q has no zeros then there is a unique B-valued regular function b on Y such that the values of the function $\widetilde{q}$ defined by the relation $b^{-1}(f\partial_z + q)b = f\partial_z + \widetilde{q}$ belong to $y + V$. This is proved just as in the case $f = 1$. Then one has to find the "normal form" of the operator $\partial_z + f^{-1}\widetilde{q}$, i.e., one must find a B-valued regular function g on $Y\backslash D$ such that $g^{-1}(\partial_z + f^{-1}\widetilde{q})g = \partial_z + \overline{q}$ where the values of $\overline{q}$ belong to $y + V$. It is easy to see that g is the image of

$$\begin{pmatrix} f & \partial_z(f) \\ 0 & 1 \end{pmatrix}$$

under the homomorphism $PGL(2) \to G$ corresponding to $\{h, x, y\}$. Therefore the components $\overline{q}_k$ and $\widetilde{q}_k$ with respect to the grading $\mathfrak{g} = \underset{k}{\oplus}\mathfrak{g}_k$ are related by

$$\overline{q}_k = f^{-k-1}\widetilde{q}_k, \quad k > 1$$

$$\overline{q}_1 = f^{-2}\Big(q_1 + \big(\tfrac{1}{2}\partial_z^2(f)\cdot f - \tfrac{1}{4}\big(\partial_z(f)\big)^2\big)z\Big)$$

So for any $k \geq 1$ the set of all possible $\overline{q}_k$ is precisely $\mathfrak{g}_k \otimes H^0\Big(Y, \mathcal{O}_Y\big((k+1)D\big)\Big)$. ■

Usually we will consider $\mathrm{Op}_{\mathfrak{g},D}(Y)$ as a subset of $\mathrm{Op}_{\mathfrak{g}}(Y\backslash D)$.

©.4.3 Let D and D' be finite subschemes of Y such that $D \subset D'$. We will construct a mapping $\mathrm{Op}_{\mathfrak{g},D}(Y) \to \mathrm{Op}_{\mathfrak{g},D'}(Y)$ such that the diagram

$$\begin{array}{ccc} \mathrm{Op}_{\mathfrak{g},D}(Y) & \longrightarrow & \mathrm{Op}_{\mathfrak{g},D'}(Y) \\ \big\downarrow\wr & & \big\downarrow\wr \\ \underline{\mathrm{Op}}_{\mathfrak{g},D}(Y) & \subset & \underline{\mathrm{Op}}_{\mathfrak{g},D'}(Y) \end{array} \qquad (©.33)$$

is commutative.

First of all to a B-bundle $\mathcal{F}$ and a finite subscheme $\Delta \subset X$ we will associate a new B-bundle $\mathcal{F}^{\Delta}$. Let $U \subset Y$ be an open subset containing Δ such that the restriction $\mathcal{F}_U$ of $\mathcal{F}$ to U is trivial and there is an $f \in H^0(U, \mathcal{O}_U)$ such that $\Delta \subset U$ is defined by the equation $f = 0$. Let $\lambda : \mathbb{G}_m \to H$ denote the homomorphism such that for any simple root α $\lambda(t)$ acts on $\mathfrak{g}^{\alpha}$ as multiplication by t. Fix a trivialization $\tau : \mathcal{F}_U \xrightarrow{\sim} U \times B$. Let us explain that by a B-bundle we understand a principle B-bundle, e.g., $\mathcal{F}_U$ is a space with a free right action of B such that $\mathcal{F}_U/B = U$. So the group of automorphisms of the B-bundle $U \times B$ is the group of regular functions $U \to B$ which acts on $U \times B$ by left multiplication.

Definition. *The B-bundle $\mathcal{F}^{\Delta}$ is obtained by gluing together $\mathcal{F}_{Y\backslash\Delta}$ and $U \times B$ by means of the composition $\mathcal{F}_{U\backslash\Delta} \xrightarrow{\tau} (U\backslash\Delta) \times B \xrightarrow{\lambda(f)} (U\backslash\Delta) \times B$.*

To check that $\mathcal{F}^{\Delta}$ is well defined we must show that if $\tilde{\tau} : \mathcal{F}_U \xrightarrow{\sim} U \times B$ is another trivialization then the function $b : U\backslash\Delta \to B$ defined by $b = \big(\lambda(f)\tilde{\tau}\big)\big(\lambda(f)\tau\big)^{-1}$ is regular on U. Indeed, $b = \lambda(f)(\tilde{\tau}\tau^{-1})\lambda(f)^{-1}$, $\tilde{\tau}\tau^{-1}$ is a regular function $U \to B$, and the coweight corresponding to λ is dominant.

Remark. For a B-module V we have a canonical grading $V = \underset{k}{\oplus} V_k$ such that $\lambda(t)x = t^k x$ for $x \in V_k$. The filtration $V^k = \underset{r \geq k}{\oplus} V_r$ is B-invariant. So for any B-bundle $\mathcal{F}$ on Y it induces the filtration $V^k_{\mathcal{F}}$ of the vector bundle $V_{\mathcal{F}}$. It is easy to show that

$$V_{\mathcal{F}^{\Delta}} = \sum_k V^k_{\mathcal{F}}(k\Delta) \qquad (\copyright.34)$$

According to Theorem ©.3.2 from [DeM] a B-bundle $\mathcal{F}$ is uniquely determined by the tensor functor $V \mapsto V_{\mathcal{F}}$ from the category of finite dimensional B-modules to the category of vector bundles on Y. So (©.34) can be considered as another definition of $\mathcal{F}^{\Delta}$. ■

Let $(\mathcal{F}, \nabla)$ be a $\mathfrak{g}$-oper on Y with D-singularities and D' a finite subscheme on Y containing D. We are going to construct a $\mathfrak{g}$-oper $(\mathcal{F}', \nabla')$ on Y with D'-singularities. Set $\mathcal{F}' = \mathcal{F}^{\Delta}$ where $\Delta \subset Y$ is a subscheme such that the corresponding ideal $\mathcal{J}_{\Delta} \subset \mathcal{O}_Y$ is equal to $\mathcal{J}_{D'}(D)$ (in other words if Δ, D, and D' are considered as effective divisors on Y then $\Delta = D' - D$). Let $\mathcal{E}_{\mathcal{F}}$ and $\mathcal{E}^{-1}_{\mathcal{F}}$ have the same meaning as in ©.1.1. $\mathcal{F}'$ and $\mathcal{F}$ are canonically isomorphic outside Δ. So we have a canonical isomorphism between the restrictions of $\mathcal{E}^{-1}_{\mathcal{F}}$ and $\mathcal{E}^{-1}_{\mathcal{F}'}$ to $Y\backslash D$. It is easy to show that it is induced by a morphism $\mathcal{E}^{-1}_{\mathcal{F}} \to \mathcal{E}^{-1}_{\mathcal{F}'}(\Delta)$ such

that the corresponding mapping $\mathcal{E}_{\mathcal{F}}^{-1}/\mathcal{E}_{\mathcal{F}} \to \left(\mathcal{E}_{\mathcal{F}'}^{-1}/\mathcal{E}_{\mathcal{F}'}\right)(\Delta)$ is an isomorphism. We define $\nabla' : \Theta(-D') \to \mathcal{E}_{\mathcal{F}'}^{-1}$ to be the composition $\Theta(-D') \xrightarrow{\nabla} \mathcal{E}_{\mathcal{F}}^{-1}(-\Delta) \to \mathcal{E}_{\mathcal{F}'}^{-1}$.

So for $D \subset D'$ we have constructed a canonical mapping $\mathrm{Op}_{\mathfrak{g},D}(Y) \to \mathrm{Op}_{\mathfrak{g},D'}(Y)$. It is easy to see that the diagram (©.33) is commutative. So the mapping $\mathrm{Op}_{\mathfrak{g},D}(Y) \to \mathrm{Op}_{\mathfrak{g},D'}(Y)$ is injective (this also follows from Proposition ©.4.2). We will usually identify $\mathrm{Op}_{\mathfrak{g},D}(Y)$ with its image in $\mathrm{Op}_{\mathfrak{g},D'}(Y)$.

©§5. The algebra $A_{\mathfrak{g},D}(X)$.

©.5.1 Let X be a smooth projective curve over $\mathbb{C}$. Then for any finite subscheme $D \subset X$ the affine space $\underline{\mathrm{Op}}_{\mathfrak{g},D}(X)$ defined in ©.4.1 is finite dimensional. So the bijection

$$\mathrm{Op}_{\mathfrak{g},D}(X) \xrightarrow{\sim} \underline{\mathrm{Op}}_{\mathfrak{g},D}(X) \tag{©.35}$$

defined in ©.4.2 induces a structure of algebraic variety on $\mathrm{Op}_{\mathfrak{g},D}(X)$. Of course this structure can be defined without using (©.35) because there is a natural notion of a family of $\mathfrak{g}$-opers with D-singularities parameterized by a scheme (the bijection (©.35) is just a convenient tool for proving that the corresponding functor $\{Schemes\} \longrightarrow \{Sets\}$ is representable by an algebraic variety isomorphic to an affine space). On the contrary, the structure of affine space on $\mathrm{Op}_{\mathfrak{g},D}(X)$ induced by (©.35) is rather "suspicious."

Denote by $A_{\mathfrak{g},D}(X)$ and $\underline{A}_{\mathfrak{g},D}(X)$ the coordinate rings of $\mathrm{Op}_{\mathfrak{g},D}(X)$ and $\underline{\mathrm{Op}}_{\mathfrak{g},D}(X)$. We have the isomorphism

$$A_{\mathfrak{g},D}(X) \xrightarrow{\sim} \underline{A}_{\mathfrak{g},D}(X) \tag{©.36}$$

induced by (©.35). Of course $A_{\mathfrak{g},D}(X)$ and $\underline{A}_{\mathfrak{g},D}(X)$ are non-canonically isomorphic to $\mathbb{C}[x_1, \ldots, x_N]$ where N can be easily computed.

The vector space $V_D^{\mathfrak{g}}(X)$ corresponding to the affine space $\mathrm{Op}_{\mathfrak{g},D}(X)$ has a canonical grading $V_D^{\mathfrak{g}}(X) = \underset{k}{\oplus} V_{D,k}^{\mathfrak{g}}(X)$ (see (©.31)). Choosing a point $P \in \underline{\mathrm{Op}}_{\mathfrak{g},D}(X)$ one obtains an isomorphism $\underline{\mathrm{Op}}_{\mathfrak{g},D}(X) \xrightarrow{\sim} V_D^{\mathfrak{g}}(X)$ and therefore a grading of $\underline{A}_{\mathfrak{g},D}(X)$. The increasing filtration $\underline{A}_{\mathfrak{g},D}^{(n)}(X)$ corresponding to this grading does not depend on P. It induces a filtration on $A_{\mathfrak{g},D}(X)$. We will give a definition of this filtration which does not involve (©.35). We will also find the corresponding graded algebra.

©.5.2 Let $h \in \mathbb{C}$ (one can consider h as "Planck's constant"). Let Y be a smooth curve over $\mathbb{C}$ and G an algebraic group over $\mathbb{C}$. If $\mathcal{F}$ is a G-bundle and $\mathcal{E}_{\mathcal{F}}$ the algebroid of infinitesimal symmetries of $\mathcal{F}$ then by an *h-connection* on $\mathcal{F}$ we mean a morphism $\nabla : \Theta \to \mathcal{E}_{\mathcal{F}}$ such that the diagram



is commutative. If $\mathcal{L}$ is a locally free sheaf of $\mathcal{O}_Y$-modules then an h-connection on $\mathcal{L}$ is a $\mathbb{C}$-linear morphism $\nabla : \mathcal{L} \to \mathcal{L} \otimes \Omega$ such that $\nabla(fs) = f\nabla(s) + hs \otimes df$ where f and s are local sections of $\mathcal{O}$ and $\mathcal{L}$.

If $h = 1$ then an h-connection is just a connection. The case $h \neq 0$ can be reduced to $h = 1$ (replace ∇ by $h^{-1}\nabla$). If $h = 0$ then an h-connection on a vector bundle $\mathcal{L}$ is an $\mathcal{O}_Y$-linear morphism $\mathcal{L} \to \mathcal{L} \otimes \Omega$ and an h-connection on a principle G-bundle $\mathcal{F}$ is an element of $H^0(Y, \mathfrak{g}_{\mathcal{F}} \otimes \Omega)$ where $\mathfrak{g}_{\mathcal{F}}$ is the vector bundle corresponding to $\mathcal{F}$ and the adjoint representation of G.

Now let G be a connected reductive group over $\mathbb{C}$. Replacing the word "connection" in Definition ©.1.2 by "h-connection" one obtains the notion of (G, h)-*oper*. In the same way one introduces the notions of $(\mathfrak{g}, h)$-*oper* and $(\mathfrak{g}, h)$-*oper with D-singularities.* Denote by $\mathrm{Op}^h_{\mathfrak{g}}(Y)$ $\big($resp. $\mathrm{Op}^h_{\mathfrak{g},D}(Y)\big)$ the set of $(\mathfrak{g}, h)$-opers on Y (resp. $(\mathfrak{g}, h)$-opers on Y with D-singularities).

Most of the results of ©§1–©§4 concerning G-opers, $\mathfrak{g}$-opers and $\mathfrak{g}$-opers with D-singularities are also true for (G, h)-opers, $(\mathfrak{g}, h)$-opers, and $(\mathfrak{g}, h)$-opers with D-singularities (the only exception is the interpretation of analytic $sl(2)$-opers as locally projective structures in ©.2.7). Of course one should replace connections by h-connections and differential operators by *h-differential operators*.

By definition, the sheaf of h-differential operators $\mathcal{D}_h$ is the sheaf of rings generated by the ring $\mathcal{O}$ and the left $\mathcal{O}$-module Θ with the following defining relations: if $v \in \Theta$ (i.e., v is a local section of Θ) and $f \in \mathcal{O}$ then $vf = fv + hv(f)$ where $v(f)$ is the derivative of f with respect to v (if $\dim Y$ were greater than 1 we would also have to add the relations $v_1v_2 - v_2v_1 = h[v_1, v_2]$ for $v_1, v_2 \in \Theta$ but if $\dim Y = 1$ these relations hold automatically). If $h = 1$ then $\mathcal{D}_h = \mathcal{D}$ while if $h = 0$ then $\mathcal{D}_h$ is the symmetric algebra of Θ. For any h there

is a canonical morphism $\mathcal{D}_h \to \mathcal{D}$ which is identical on $\mathcal{O}$ and acts on Θ as multiplication by h. It is an isomorphism for $h \neq 0$.

In ©.2.6 we associated to each $sl(2)$-oper a differential operator $\Omega^{\otimes(-1/2)} \to \Omega^{\otimes(3/2)}$ of order 2. In the same way one associates to an $(sl(2), h)$-oper an h-differential operator $\Omega^{\otimes(-1/2)} \to \Omega^{\otimes(3/2)}$. Applying the canonical morphism $\mathcal{D}_h \to \mathcal{D}$ one obtains a differential operator $\Omega^{\otimes(-1/2)} \to \Omega^{\otimes(3/2)}$ with principal symbol h^2. Thus we get a natural bijection between $\mathrm{Op}^h_{sl(2)}(Y)$ and the set of differential operators $L : \Omega^{\otimes(-1/2)} \to \Omega^{\otimes(3/2)}$ of order 2 such that $L^t = L$ and the principal symbol of L equals h^2. In particular $\mathrm{Op}^h_{sl(2)}(Y)$ is a principal homogenous space over $H^0\big(Y, \Omega^{\otimes 2}\big)$. The description of the action of $H^0\big(Y, \Omega^{\otimes 2}\big)$ on $\mathrm{Op}_{sl(2)}(Y)$ given in ©.3.2 works also for $\mathrm{Op}^h_{sl(2)}(Y)$.

Now let $\mathfrak{g}$ be an arbitrary semisimple Lie algebra over $\mathbb{C}$ and $V^{\mathfrak{g}}_D(Y)$ the vector space defined by (©.31). Denote by $\underline{\mathrm{Op}}^h_{\mathfrak{g},D}(Y)$ the principal homogeneous space over $V^{\mathfrak{g}}_D(Y)$ induced by $\mathrm{Op}^h_{sl(2)}(Y)$ via the embedding $H^0\big(Y, \Omega^{\otimes 2}\big) \hookrightarrow V^{\mathfrak{g}}_D(Y)$ defined by (©.26). Just as in ©.3.4 and ©.4.2 one constructs a canonical bijection

$$\underline{\mathrm{Op}}^h_{\mathfrak{g},D}(Y) \xrightarrow{\sim} \mathrm{Op}^h_{\mathfrak{g},D}(Y) \qquad (©.37)$$

©.5.3 Let X be a smooth projective curve over $\mathbb{C}$ and D a finite subscheme of X. Set

$$\mathcal{OP}_{\mathfrak{g},D}(X) = \Big\{(h,u) \;\Big|\; h \in \mathbb{C},\ u \in \mathrm{Op}^h_{\mathfrak{g},D}(X)\Big\}$$
$$\underline{\mathcal{OP}}_{\mathfrak{g},D}(X) = \Big\{(h,u) \;\Big|\; h \in \mathbb{C},\ u \in \underline{\mathrm{Op}}^h_{\mathfrak{g},D}(X)\Big\}$$

There is a natural structure of algebraic variety on $\mathcal{OP}_{\mathfrak{g},D}(X)$ and $\underline{\mathcal{OP}}_{\mathfrak{g},D}(X)$ and (©.37) induces an isomorphism $\underline{\mathcal{OP}}_{\mathfrak{g},D}(X) \xrightarrow{\sim} \mathcal{OP}_{\mathfrak{g},D}(X)$. So $\mathcal{OP}_{\mathfrak{g},D}(X)$ is an affine variety non-canonically isomorphic to an affine space and the morphism $\pi : \mathcal{OP}_{\mathfrak{g},D}(X) \to \mathbb{A}^1$ which maps $\mathrm{Op}^h_{\mathfrak{g},D}(X)$ to $h \in \mathbb{C}$ is flat. $\mathbb{G}_m$ acts on $\mathcal{OP}_{\mathfrak{g},D}(X)$ (replacing ∇ by $\lambda\nabla$, $\lambda \in \mathbb{C}^*$) and π is $\mathbb{G}_m$-equivariant. Set $\mathcal{OP}^*_{\mathfrak{g},D}(X) = \pi^{-1}(\mathbb{G}_m)$. We have $\mathrm{Op}_{\mathfrak{g},D}(X) = \pi^{-1}(1) = \mathbb{G}_m \backslash \mathcal{OP}^*_{\mathfrak{g},D}(X)$. Therefore $A_{\mathfrak{g},D}(X)$, the coordinate ring of $\mathrm{Op}_{\mathfrak{g},D}(X)$, can be identified with the ring of $\mathbb{G}_m$-invariant regular functions on $\mathcal{OP}^*_{\mathfrak{g},D}(X)$. So we have a filtration $A_{\mathfrak{g},D}(X) = \bigcup\limits_n A^{(n)}_{\mathfrak{g},D}(X)$ where $A^{(n)}_{\mathfrak{g},D}(X)$ is the space of $\mathbb{G}_m$-invariant regular functions on $\mathcal{OP}^*_{\mathfrak{g},D}(X)$ having a pole of order $\leq n$ at the divisor $\pi^{-1}(0)$. It is easy to show that this filtration corresponds to the filtration $\underline{A}_{\mathfrak{g},D}(X) = \bigcup\limits_n \underline{A}^{(n)}_{\mathfrak{g},D}(X)$ defined in ©.5.1.

©.5.4 The graded ring $grA_{\mathfrak{g},D}(X) = \underset{n}{\oplus} A^{(n)}_{\mathfrak{g},D}(X)/A^{(n-1)}_{\mathfrak{g},D}(X)$ is nothing but the coordinate ring of $\pi^{-1}(0) = \mathrm{Op}^0_{\mathfrak{g},D}(X)$. The grading on the coordinate ring comes from the action of $\mathbb{G}_m$ on $\mathrm{Op}^0_{\mathfrak{g},D}(X)$. We will describe $\mathrm{Op}^0_{\mathfrak{g},D}(X)$ together with this action. In what follows we use the notation of ©.1.1 and ©.1.2.

Let G be the connected algebraic group with trivial center corresponding to $\mathfrak{g}$, B its Borel subgroup. By definition, points of $\mathrm{Op}^0_{\mathfrak{g},D}(X)$ correspond to pairs $(\mathcal{F}, \omega)$ where $\mathcal{F}$ is a B-bundle on X and $\omega \in H^0\big(X, \mathfrak{g}^{-1}_{\mathcal{F}} \otimes \Omega(D)\big)$ is such that for every simple root α the image of ω in $H^0\big(X, \mathfrak{g}^{-\alpha}_{\mathcal{F}} \otimes \Omega(D)\big)$ has no zeros as a section of $\mathfrak{g}^{-\alpha}_{\mathcal{F}} \otimes \Omega(D)$. Denote by $\mathrm{Inv}(\mathfrak{g})$ the algebra of G-invariant polynomials on $\mathfrak{g}$. If $f \in \mathrm{Inv}(\mathfrak{g})$ is homogeneous of degree n then $f(\omega) \in H^0\big(X, \Omega^{\otimes n}(nD)\big)$. So we obtain a mapping

$$\mathrm{Op}^0_{\mathfrak{g},D}(X) \to \mathrm{Hitch}_{\mathfrak{g},D}(X) = \mathrm{Mor}(\mathrm{Inv}(\mathfrak{g}), \underset{n}{\oplus} H^0\big(X, \Omega^{\otimes n}(nD)\big) \qquad (©.38)$$

where Mor denotes the set of graded algebra morphisms. There is a canonical isomorphism

$$\mathrm{Inv}(\mathfrak{g}) \xrightarrow{\sim} \mathrm{Inv}(\mathfrak{g}^{\wedge *}) \qquad (©.39)$$

where $\mathrm{Inv}(\mathfrak{g}^{\wedge *})$ is the algebra of invariant polynomials of the Langlands dual $\mathfrak{g}^{\wedge}$ (indeed, both $\mathrm{Inv}(\mathfrak{g})$ and $\mathrm{Inv}(\mathfrak{g}^{\wedge *})$ are canonically isomorphic to the algebra of polynomials on $\mathfrak{h}$ invariant with respect to the Weyl group). So the r.h.s. of (©.38) can be identified with $\mathrm{Hitch}_{\mathfrak{g}^{\wedge},D}(X)$, the base of Hitchin's fibration (see 0.2), and (©.38) can be interpreted as a mapping

$$\mathrm{Op}^0_{\mathfrak{g},D}(X) \to \mathrm{Hitch}_{\mathfrak{g}^{\wedge},D}(X) \qquad (©.40)$$

$\mathrm{Hitch}_{\mathfrak{g}^{\wedge},D}(X)$ has a natural structure of affine algebraic variety with an action of $\mathbb{G}_m$ on it (see section 0).

Proposition. *The mapping (©.40) is a $\mathbb{G}_m$-equivariant isomorphism of algebraic varieties.*

Corollary. *$grA_{\mathfrak{g},D}(X)$ is canonically isomorphic to the coordinate ring of* $\mathrm{Hitch}_{\mathfrak{g},D}(X)$.

The proof of the proposition is easy: it is clear that (©.40) is a $\mathbb{G}_m$-equivariant morphism and the assertion that it is an isomorphism is an immediate consequence of the following theorem proved in [Ko2].

Kostant's Theorem. *Set* $\mathfrak{g}_*^{-1} = \Big\{u + \sum\limits_{\alpha\in\Gamma} v_\alpha \;\Big|\; u \in \mathfrak{b},\ v_\alpha \in \mathfrak{g}^{-\alpha},\ v_\alpha \neq 0\Big\}$ *where* Γ *is the set of simple roots. Denote by* f *the canonical mapping* $\mathfrak{g} \to \operatorname{Spec}\operatorname{Inv}(\mathfrak{g}) = W\backslash\mathfrak{h}$ *where* W *is the Weyl group. Then*

(i) there is an affine subspace $L \subset \mathfrak{g}_*^{-1}$ *such that the mapping* $B \times L \to \mathfrak{g}_*^{-1}$, $(b, v) \mapsto bvb^{-1}$, *is an isomorphism of algebraic varieties*

(ii) the mapping $B\backslash\mathfrak{g}_*^{-1} = L \to W\backslash\mathfrak{h}$ *induced by* f *is an isomorphism of algebraic varieties.*

For the reader's convenience we sketch a proof of Kostant's theorem. Let $\{h, x, y\}$ be the $sl(2)$-triple considered in ©.3.4. Set $V = \operatorname{Ker}\operatorname{ad} x$ and $L = y + V$. The proof of the statement (i) is quite similar to that of Proposition ©.1.3 and Theorem ©.3.4 (one uses the equalities $\mathfrak{g} = (\operatorname{Im}\operatorname{ad} y) \oplus V$ and $(\operatorname{Ker}\operatorname{ad} y) \cap \mathfrak{b} = 0$). The mapping $\mathfrak{h} \to B\backslash\mathfrak{g}_*^{-1}$ which associates to $a \in \mathfrak{h}$ the b-conjugacy class of $a + y$ factors through $W\backslash\mathfrak{h}$ because for every simple root α

$$s_\alpha(a) + y = \exp\big(-\alpha(a) \cdot \operatorname{ad} u_\alpha\big)(a + y), \quad a \in \mathfrak{h}$$

where $s_\alpha(a) = a - \alpha(a)\check{\alpha}$ and $u_\alpha \in \mathfrak{g}^\alpha$ is such that $[u_\alpha, y] = \check{\alpha}$. The composition $W\backslash\mathfrak{h} \to B\backslash\mathfrak{g}_*^{-1} \to W\backslash\mathfrak{h}$ is identical because the closure of the conjugacy class of $a + y$ contains a for $a \in \mathfrak{h}$. It remains to show that $\dim(B\backslash\mathfrak{g}_*^{-1}) = \dim(W\backslash\mathfrak{h})$, i.e., $\dim V = \dim\mathfrak{h}$. Indeed, $\dim V = \dim\operatorname{Ker}\operatorname{ad} x = n$ where n is the number of irreducible components of $\mathfrak{g}$ considered as a module over $sl(2) = \mathbb{C}h + \mathbb{C}x + \mathbb{C}y$. But since the eigenvalues of $\operatorname{ad} h$ are even $n = \dim\operatorname{Ker}\operatorname{ad} h = \dim h$.

Remark. There is a canonical $\mathbb{G}_m$-invariant isomorphism of algebraic varieties

$$V_D^{\mathfrak{g}}(X) \xrightarrow{\sim} \operatorname{Hitch}_{\mathfrak{g}^\wedge, D}(X) \tag{©.41}$$

where $V_D^{\mathfrak{g}}(X)$ is defined by (©.31) and $\lambda \in \mathbb{G}_m$ acts on $V_{D,k}^{\mathfrak{g}}(X)$ as multiplication by λ^k. To construct (©.41) choose a principal $sl(2)$-triple $\{h, x, y\}$ and identify the spaces $V_{k-1}(\mathfrak{g})$ from (©.31)) with the eigenspaces of $\operatorname{ad} h$ in $\operatorname{Ker}\operatorname{ad} x$. Then to every $\eta \in V_D^{\mathfrak{g}}(X)$ and every homogeneous $f \in \operatorname{Inv}(\mathfrak{g})$ of degree n we associate $f(y + \eta) \in H^0\big(X, \Omega^{\otimes n}(nD)\big)$. This defines the mapping (©.41) and Kostant's theorem shows that it is an isomorphism.

References

[De1] P. Deligne, *Équations differentielles à points singuliers réguliers*, Lecture Notes in Mathematics, Springer-Verlag, Berlin, vol. **163** (1970).

[DeM] P. Deligne and J. Milne, *Tannakian categories*, Lecture Notes in Mathematics, Springer-Verlag, Berlin, vol. **900** (1982), p. 101–228.

[DS1] V. G. Drinfeld and V. V. Sokolov, *Equations of Korteweg-deVries type and simple Lie algebras*, Soviet Mathematics Doklady, vol. **23** (1981), No. 3, p. 457–462.

[DS2] V. G. Drinfeld and V. V. Sokolov, *Lie algebras and equations of Korteweg-deVries type*, Journal of Soviet Mathematics, vol. **30** (1985), p. 1975–2035.

[Hi1] N. J. Hitchin, *Stable bundles and integrable systems*, Duke Math. Journal, vol. **54** (1987), p. 91–114.

[Ko1] B. Kostant, *The principal three-dimensional subgroup and the Betti numbers of a complex simple Lie group*, American Journal of Math., vol. **81** (1959), p. 973–1032.

[Ko2] B. Kostant, *Lie group representations on polynomial rings*, American Journal of Math., vol. **85** No. 3, (1963), p. 327–404.

[Te1] C. Teleman, *Sur les structures homographiques d'une surface de Riemann*, Comment. Math. Helv., vol. **33** (1959), p. 206–211.

[Tyu] A. N. Tyurin, *On the periods of quadratic differentials*, Uspekhi Mat. Nauk, vol. **33** No. 6, (1978), p. 149–195 (Russian).